

Resistance to Anti-EGFR Therapy and Strategies to Overcome it: Possible Role of BDNF/TrkB

de Farias CB^{1,2*}, Schwartzmann G^{1,3} and Roesler R^{1,4}

¹Cancer and Neurobiology Laboratory, Clinical Hospital, Federal University of Rio Grande do Sul, RS, Brazil

²Children's Cancer Institute, Porto Alegre, RS, Brazil

³Department of Internal Medicine, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil

⁴Department of Pharmacology, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil

Abstract

Acquired resistance to anti-epidermal growth factor receptor (EGFR) monoclonal antibodies such as cetuximab and panitumumab represents an important limitation to the treatment of colorectal cancer (CRC), and research efforts have increasingly been directed towards understanding molecular mechanisms of resistance. We propose that neurotrophin signaling mediated by brain-derived neurotrophic factor (BDNF) and its receptor, tropomyosin-related kinase B (TrkB), might be a compensatory mechanism contributing to decreased response to anti-EGFR therapy in CRC. The combined targeting of EGFR and TrkB is a novel strategy worth of further investigation.

Keywords: Brain-derived neurotrophic factor; TrkB; Neurotrophin; Epidermal growth factor receptor; Cetuximab; Colorectal cancer

Short Communication

The emergence of secondary resistance to monoclonal antibodies that selectively target the epidermal growth factor receptor (EGFR), cetuximab and panitumumab, constitutes a current challenge in the treatment of patients with colorectal cancer (CRC). Although most research has focused particularly on genetic alterations that activate downstream signaling effectors, such as mutations in *KRAS*, *BRAF*, and *PIK3CA* [1], the contribution to resistance of compensation by parallel tyrosine kinase receptor pathways has also been recently highlighted [2].

In this context, we propose that an emerging target is the tropomyosin-related kinase B (TrkB, also called tropomyosin receptor kinase B or receptor tyrosine kinase B), encoded by *NTRK2* and activated by the neurotrophin brain-derived neurotrophic factor (BDNF). TrkB shares several downstream effectors with EGFR, including the mitogen-activated protein kinase (MAPK) and phosphatidylinositol 3-kinase (*PI3K*) pathways, and activation of EGFR plays a role in upregulating BDNF mRNA and protein levels in the brain [3]. In neurons, BDNF/TrkB signaling plays an important role for normal development, survival and plasticity [4], but it has also been proposed to play a role in cancer progression, metastasis, and resistance to chemotherapy [5].

We and others have reported that BDNF and TrkB levels are increased in CRC tumors [6,7]. Treating human CRC cells with an experimental anticancer peptide increases the mRNA expression, protein content, and secretion of BDNF, through a mechanism that depends on EGFR activity. The increase in BDNF results in a compensatory response that rescues CRC cell proliferation and survival [6]. Importantly, BDNF protects against the anti-proliferative effect of cetuximab in a human CRC cell line, whereas a TrkB inhibitor can sensitize the cells to a low dose of cetuximab [8].

This evidence raises the possibility that BDNF/TrkB upregulation might contribute to resistance against anti-EGFR therapy in CRC. Whether tumors from patients with intrinsic resistance to anti-EGFR therapy present increased BDNF and TrkB expression and activity remains to be determined. It is possible that the combined inhibition of EGFR and TrkB could increase tumor responses to treatment [8,9].

This may be a novel opportunity to overcome resistance to anti-EGFR therapy, and it warrants further experimental exploration.

Acknowledgements

The authors are supported by the National Council for Scientific and Technological Development (CNPq; grant numbers 484185/2012-8 and 303276/2013-4 to R.R., and 309430/2012-7 to G.S.); the Clinical Hospital research fund (FIPE-HCPA); the South American Office for Anticancer Drug Development; and the Children's Cancer Institute (ICI).

References

1. Bardelli A, Siena S (2010) Molecular mechanisms of resistance to cetuximab and panitumumab in colorectal cancer. *J Clin Oncol* 28: 1254-1261.
2. Bertotti A, Sassi F (2015) Molecular pathways: sensitivity and resistance to anti-EGFR antibodies. *Clin Cancer Res* 21: 3377-3383.
3. Tirassa P, Triaca V, Amendola T, Fiore M, Aloe L (2003) EGF and NGF injected into the brain of old mice enhance BDNF and ChAT in proliferating subventricular zone. *J Neurosci Res* 72: 557-564.
4. Huang EJ, Reichardt LF (2003) Trk receptors: roles in neuronal signal transduction. *Annu Rev Biochem* 72: 609-642.
5. Thiele CJ, Li Z, McKee AE (2009) On Trk-the TrkB signal transduction pathway is an increasingly important target in cancer biology. *Clin Cancer Res* 15: 5962-5967.
6. Brunetto de Farias C, Rosemberg DB, Heinen TE, Koehler-Santos P, Abujamra AL, et al. (2010) BDNF/TrkB content and interaction with gastrin-releasing peptide receptor blockade in colorectal cancer. *Oncology* 79: 430-439.
7. Dawson H, Grundmann S, Koelzer VH, Galvan JA, Kirsch R, et al. (2015) Tyrosine kinase receptor B (TrkB) expression in colorectal cancers highlights anoikis resistance as a survival mechanism of tumour budding cells. *Histopathology* 66: 715-725.

***Corresponding author:** de Farias CB, Cancer and Neurobiology Laboratory, Experimental Research Center, Clinical Hospital, Federal University of Rio Grande do Sul. Rua Ramiro Barcelos, 2350, CPE, 2^a Andar, 90035-003 Porto Alegre, RS, Brazil, Tel: +555151 33597616; Fax: +5551 33598012; E-mail: labpesquisa@ci-rs.org.br

Received December 30, 2015; **Accepted** February 24, 2016; **Published** February 26, 2016

Citation: de Farias CB, Schwartzmann G, Roesler R (2016) Resistance to Anti-EGFR Therapy and Strategies to Overcome it: Possible Role of BDNF/TrkB. *J Cancer Sci Ther* 8: 046-047. doi:10.4172/1948-5956.1000387

Copyright: © 2016 de Farias CB, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

8. de Farias CB, Heinen TE, dos Santos RP, Abujamra AL, Schwartzmann G, et al. (2012) BDNF/TrkB signaling protects HT-29 human colon cancer cells from EGFR inhibition. *Biochem Biophys Res Commun* 425: 328-332.
9. Roesler R, de Farias CB, Abujamra AL, Brunetto AL, Schwartzmann G (2011) BDNF/TrkB signaling as an anti-tumor target. *Expert Rev Anticancer Ther* 11: 1473-1475.