

Discussiones Mathematicae

Probability and Statistics 29 (2009) 155–168

doi:10.7151/dmps.1112

A DOMINANT HEIGHT GROWTH MODEL FOR EUCALYPTUS PLANTATIONS IN PORTUGAL

AYANA MATEUS

*CMA, Departamento de Matemática,
Faculdade de Ciências e Tecnologia FCT
Universidade Nova de Lisboa, UNL
Quinta da Torre, 2829–516 Caparica, Portugal*

e-mail: amf@fct.unl.pt ; +351 21 2948388

AND

MARGARIDA TOMÉ

*CEF-Centro de Estudos florestais
ISA- Instituto Superior de Agronomia
UTLisboa-Universidade Técnica de Lisboa
Tapada da Ajuda, 1349–017 Lisboa, Portugal*

e-mail: magatome@isa.utl.pt ; +351 21 3653100

Abstract

Eucalyptus globulus Labill is one of the most important economic forest species in Portugal, occupying an area of $875.10^3 ha$ in a total forest area of $3346.10^3 ha$ (Tomé *et al.*, 2007). The main goal of this study is to develop a dominant height growth model for Eucalyptus, applicable throughout the country, representing an improve of the curves that are part of the whole stand model existing in Portugal, the GLOBULUS model (Tomé *et al.*, 2001). The dominant height growth model will be built on a biological function formulated as a difference equation (Algebraic Difference Approach) to an all possible growth intervals data structure. The dynamic model proposed, obtains non biased and efficient estimates for the parameters. Comparing to the

already available model, the last has the advantage of expressing the asymptote in weather variables functions, meaning that it is possible to reproduce the eucalyptus growth according to weather changes.

Keywords: algebraic difference equation, dynamic model, dominant height, forest planning, non linear model.

2000 Mathematics Subject Classification: Primary 62J02; Secondary 62P12.

REFERENCES

- [1] M. Tomé, S. Barreiro, A. Cortiçada, A. Meyer, T. Ramos and P. Malico, Inventário florestal 2005-2006. Áreas, volumes e biomassas dos povoamentos florestais. Resultados Nacionais e por Nut's II e III. Publicações GIMREF RT 5/2007 Universidade Técnica de Lisboa. Instituto Superior de Agronomia, Centro de Estudos Florestais, Lisboa, Portugal 2007.
- [2] M. Tomé, F. Ribeiro and P. Soares, O modelo GLOBULUS 2.1-Relatórios técnico científicos do GIMREF nº 1/2001. Universidade Técnica de Lisboa, Instituto Superior de Agronomia, Centro de Estudos Florestais, Lisboa, Portugal 2001.
- [3] G.G. Wang, K. Klinka and W. John, *Use of synoptic variables in predicting white spruce site index*, For. Ecol. Manage **80** (1996), 95–105.
- [4] R.A. Monserud, *Height growth and site index curves for inland Douglas-fir based on stem analysis data and forest habitat type*, For. Sci. **30** (1984), 943–965.
- [5] L. Clutter, J. Fortson, L. Pienaar, G. Brister and L. Bailey, *Timber Management: A Quantitative approach*, John Wiley and Sons 1983.
- [6] F. Ribeiro and M. Tomé, *Classificação climática de Portugal continental, baseada em informação do atlas do ambiente*, Revista de Ciências Agrárias 2000.
- [7] A.R. Gallant, *Nonlinear Statistical Models*, John Wiley & Sons (1987), 624p.
- [8] The SAS System for windows, 9.1 edition.
- [9] G.A.F. Seber and C.G. Wild, *Nonlinear Regression*, New York, John Wiley 1989.

Received 17 September 2009