



## ACUTE PHASE PROTEINS IN SHEEP AND GOATS – FUNCTION, REFERENCE RANGES AND ASSESSMENT METHODS: AN OVERVIEW

P. T. ILIEV<sup>1</sup> & T. M. GEORGIEVA<sup>2</sup>

<sup>1</sup>Department of Veterinary Microbiology, Infectious and Parasitic Diseases, <sup>2</sup>Department of Pharmacology, Animal Physiology and Physiological Chemistry, Faculty of Veterinary medicine, Trakia University, Stara Zagora, Bulgaria

### Summary

Iliev, P. T. & T. M. Georgieva, 2018. Acute phase proteins in sheep and goats – function, reference ranges and assessment methods: An overview. *Bulg. J. Vet. Med.*, **21**, No 1, 1–16.

Acute phase response (APR) is activated by disorders in systemic homeostasis. The main purpose of APR is to prevent further tissue damage by isolation and destruction of causative agent, removing the toxic products and providing conditions for tissue repair. Probably the most significant change during APR is a transformation in the liver protein spectrum expressed by increased rate of synthesis of acute phase proteins (APPs) or acute phase reactants. Numerous APPs that perform specific functions are established whose quantitative variations vary considerably among different animals. Some of them are mediators (C-reactive protein, fibrinogen), others act as inhibitors (protease inhibitors –  $\alpha$ 1-antitrypsin) and especially in small ruminants, the most specific APPs are scavengers and transporters (haptoglobin, serum amyloid A). Despite considerable progress, many of the characteristics of APPs in domestic animals are still poorly understood especially in sheep and goats. Little is known about some other proteins such as lipopolysaccharide binding protein, hemopexin, alpha-1 antitrypsin, lactoferrin, transferrin, C-reactive protein, ceruloplasmin, fibrinogen and alpha-1 acid glycoprotein. The aim of this study is to present information concerning the most important functions of APPs, as well as their ranges in healthy sheep and goats and laboratory assay methods.

**Key words:** acute phase proteins, sheep, goats

### INTRODUCTION

The acute phase response (APR) or acute phase reaction is a complex early, antigen non-specific, defense systemic reaction appearing before a specific immune response is raised (Petersen *et al.*, 2004). It is activated by trauma, neoplastic growth, bacterial, parasitic and viral infection, burns, surgery, immunological disorders

etc. (Gruys *et al.*, 2005; Cray *et al.*, 2009; Eckersall & Bell, 2010; Tothova *et al.*, 2014). The main purpose of APR is to restore the homeostasis by isolating and destroying the harmful agent and to activate the repair process (Janeway *et al.*, 2001; Ceciliani *et al.*, 2002). The damaged tissue is the primary source that initi-

ates APR. The early changes involve activation of cells belonging to the innate immunity – macrophages and monocytes which synthesise pro-inflammatory cytokines such as interleukin-1 (IL-1), interleukin-6 (IL-6), tumour necrosis factor alpha (TNF- $\alpha$ ) and some others (Murata *et al.*, 2004). These mediators are involved in the activation of fibroblasts, leukocytes and endothelial cells. The latter, by producing more cytokines (IL-6, IL-8, IL-11, PGE<sub>2</sub>, LIF, nitric oxide), increase the permeability of blood vessels, enhance leukocyte migration and induce local changes – swelling, pain, redness and warmth which accompany the inflammatory process (Gruys *et al.*, 2005). The increased concentration of cytokines may induce systemic APR and resultant activation of the hypothalamic-pituitary-adrenal system (increased adrenocorticotrophic hormone and glucocorticoids production), reduction of growth hormone secretion and a considerable number of metabolic and biochemical changes (complement cascade and blood coagulation system activation; decreased serum concentration of iron, zinc, calcium, vitamin A,  $\alpha$ -tocopherol; negative nitrogen balance; changes in the concentration of some plasma proteins) clinically manifested by pyrexia, anorexia, weight loss etc. (Moshage, 1997; Gabay & Kushner, 1999; Gruys *et al.*, 1999; Pyorala, 2000; Gruys *et al.*, 2005; Jain *et al.*, 2011; Ceciliani *et al.*, 2012; O'Reilly & Eckersall, 2014; Tothova *et al.*, 2014). Down-regulation of APR is controlled by many inflammatory mediators (IL-4, IL-10) and hormones (glucocorticoids) as well as by the production of antagonists to pro-inflammatory cytokines (Gruys *et al.*, 2005). Usually APR continues for 1–2 days and after this period the affected organism should return to normal function

(Ceciliani *et al.*, 2002). If the acute inflammation becomes chronic, APR is still induced (Baumann & Gauldie, 1994; Jain *et al.*, 2011).

One of the most important metabolic alterations accompanying APR occurs mainly in the liver but also in other organs. It involves the synthesis of a large group proteins, collectively known as acute phase proteins (Ceciliani *et al.*, 2012). Their expression depends on cytokines' levels, whereas the cytokine activity modulation is regulated by glucocorticoids and growth factors (van Miert, 1995; Ceciliani *et al.*, 2002; Gruys *et al.*, 2005).

#### ACUTE PHASE PROTEINS

The acute phase proteins are a large group of plasma proteins which originate mainly from the liver in response to pro-inflammatory cytokines (IL-1, IL-6, TNF- $\alpha$ ). They are released very quickly into the circulation, where each performs a specific activity. During APR the serum of APPs concentrations increase substantially while in healthy animals and humans, APPs are undetectable or in negligible quantities. However, some APPs collectively called “permanent” are continuously secreted and released into the bloodstream while the “induced” ones are present in plasma only during APR. In general, the main function of APPs is to defend the host against pathological damages, assist in the restoration of homeostasis and in the regulation of different stages of inflammation (Petersen *et al.*, 2004; Tothova *et al.*, 2014).

APPs can also be synthesised extra-hepatically in some tissues such as testicular tissue, adipose tissue, lung, ovary, uterus, mammary glands, digestive tract (Ceciliani *et al.*, 2012).

The type and magnitude of APP expression varies between the species. Plasma concentrations of major APPs in healthy animals are very low. During the acute phase of inflammation their levels are greatly enhanced in the first few hours after exposure to the pathogenic factors (Gruys *et al.*, 1999; Ceciliani *et al.*, 2002; Petersen *et al.*, 2004). APPs kinetics depends on the animal species and extent of tissue damage. Serum APPs concentrations generally reach a peak within 24–48 hours. If a new stimulus does not appear and inflammation ceases, the feedback APPs regulation limits the response within 4 to 7 days after challenge. Chronic inflammation is considered as a series of individual inflammatory stimuli and is characterised by longer and slight increase in the serum concentration of APPs as compared to acute inflammation (Sipe, 1985; Murtaugh, 1994; Heegaard *et al.*, 2000; Jain *et al.*, 2011).

#### CLASSIFICATION OF ACUTE PHASE PROTEINS

Despite the uniformity in the development of APR between species, there is a lot of evidence showing that each animal has its own specific APPs set (Pyorala, 2000). The biological functions of some APPs studied in ruminants are summarised in Table 1.

In general terms, these proteins can be classified according to their quantitative variation and function.

A) Depending on the magnitude of the increase or decrease by at least 25% during APR (Murata *et al.*, 2004; Ceron *et al.*, 2005; Eckersall & Bell, 2010).

- Positive APPs – haptoglobin, serum amyloid A, C-reactive protein.
- Negative APPs – albumin, transferrin, transthyretin (prealbumin, thyroxine-binding protein).

B) Depending on the magnitude of increase in the concentration of positive

**Table 1.** Biological function of acute phase proteins (according to Ceciliani *et al.*, 2012; O'Reilly & Eckersall, 2014; Tothova *et al.*, 2014)

Proteins	Abbreviation	Biological function	Classification
Haptoglobin	Hp	Binds free haemoglobin	Positive
Serum amyloid A	SAA	Binds cholesterol, opsonin	Positive
Ceruloplasmin	Cp	Copper transport, iron metabolism	Positive
Albumin	Alb	Amino acids source, transporter, osmotic pressure	Negative
C-reactive protein	CPR	Complement activation, binding to membrane phosphorylcholine, opsonization	Positive
Fibrinogen	Fb	Blood clotting	Positive
Lipopolysaccharide binding protein	LPB	Binds to bacterial LPS, macrophage cell activation	Positive
$\alpha_1$ -acid glucoprotein	AGP	Bind drugs, inflammatory mediators and bacterial derived molecules	Positive
$\alpha_1$ -antitrypsin	AAT	Serine protease inhibitor	Positive
Transferrin	TF	Binds iron	Negative
Hemopexin	Hpx	Heme binding	Positive
Lactoferrin	LF	Binding and transferring $Fe^{3+}$ ions	Positive

APPs (Eckersall & Bell, 2010; Gomez-Laguna *et al.*, 2011; Tothova *et al.*, 2014) (Table 2).

- Major APPs – their concentrations increase 10 to 100-fold, reaching a peak 24–48 hours after pathological stimulus and decline quickly due to short half – life (Niewold *et al.*, 2003).
- Moderate APPs – their concentrations increase 5 to 10-fold, reaching a peak 2–3 days after pathological stimulus and decline more slowly than major APPs (Eckersall, 2006a).
- Minor APPs – gradually increase by between 50% and 100% of their normal level (Eckersall & Bell, 2010).

C) Depending on their role during inflammation (Tirziu, 2009; Khan & Khan, 2010).

- Mediators (C-reactive protein, fibrinogen)
- Modulators (complement proteins, inhibitors to coagulation cascade)
- Inhibitors (protease inhibitors –  $\alpha$ 1-antitrypsin,  $\alpha$ 2-macroglobulin)
- Scavengers and transporters (haptoglobin, serum amyloid A, ceruloplasmin)
- Immunomodulators ( $\alpha$ 1-acid glycoprotein)

D) Depending on their cytokine induction (Murata *et al.*, 2004; Gruys *et al.*, 2005)

- Type 1 dependent (induced by IL-1 and TNF- $\alpha$ )
- Type 2 dependent (induced by IL-6)

The synthesis of type-1 dependent APPs is synergistically affected also by IL-6, in contrast, the secretion of type-2 APPs is neither induced nor synergistically affected by IL-1, moreover it has an inhibitory effect on their synthesis. The reduction of negative APPs in plasma during inflammation is due to the priority of positive APPs synthesis. Furthermore,

some of negative APPs are hormone-binding proteins and their reduction provides more available, biologically active forms of hormones, some of which (glucocorticoids) are very important for induction of APR. Other APPs as transthyretin have also an inhibitory effect on IL-1 production by monocytes and endothelial cells, thus its declining concentration may be regarded as a part of the adaptive mechanisms of APR induction (Ceciliani *et al.*, 2002). Some APPs inhibit protease enzymes synthesised by phagocytes and pathogenic microorganisms and therefore protect tissues from damage e.g.  $\alpha$ 1-antitrypsin,  $\alpha$ 2-macroglobulin (Gruys *et al.*, 1999). Other major APPs as haptoglobin, serum amyloid A and C-reactive protein act as scavengers by binding metabolites formed during inflammation (Wagener *et al.*, 2001). Some APPs ( $\alpha$ 1-acid glycoprotein) have an antibacterial activity and thus positively influence the immune response (Eckersall, 2006b).

**Table 2.** APPs in sheep and goats

	Sheep	Goats
Major APPs	Haptoglobin, SAA	Haptoglobin, SAA
Moderate APPs	AGP	Fibrinogen, AGP
Minor APPs	Fibrinogen, Ceruloplasmin	Ceruloplasmin
Negative APPs	Albumin	Albumin

SAA= serum amyloid A; AGP= $\alpha$ 1-acid glycoprotein

## ACUTE PHASE PROTEINS IN SMALL RUMINANTS

### *Haptoglobin*

Haptoglobin (Hp) is considered to be a major APP in sheep and goats (Table 2).

It refers to a group of transporter (metal-binding) plasma proteins that increase during APR (Pannen & Robotham, 1995). Hp is synthesised mainly in the liver, although its coding gene is expressed in other tissues such as lung, skin, spleen, kidney and adipose tissue (D'Armiento *et al.*, 1997; Yang *et al.*, 2003). The synthesis of Hp is mediated and regulated by the growth hormone, insulin, bacterial endotoxin, prostaglandins, pro-inflammatory cytokines – IL-1, IL-6, tumour necrosis factor (Raynes, 1994).

Hp has several distinctive features. According to Yang *et al.* (2003) and Kato (2009) the most important one is to bind the free haemoglobin (Hb) in equimolar ratio with very high affinity (Table 1). The complex Hp-Hb is too large and cannot pass through the glomerulus of the kidney and may only be removed by reticuloendothelial system. It is accomplished by binding to CD163 receptors presented on macrophages and monocytes surface (Schaer *et al.*, 2002). Hp-Hb half-life is up to 50 min (Ceciliani *et al.*, 2012). The quantity of Hp does not increase in cases in which red blood cells are destroyed in the spleen or liver but not in the vessels (Jain *et al.*, 2011) and it may decrease during massive erythrolysis (Smith & Roberts, 1994).

The free Hb in the bloodstream is toxic and has oxidative activity (Wagner *et al.*, 2001). Hp binds it and thus prevents oxygen radicals formation (stimulated by iron), reduces the oxidative damages associated with haemolysis which defines its role as an antioxidant (Smith & Roberts, 1994; Murata *et al.*, 2004).

Hb contains iron which is one of essential elements required for bacterial growth. After the Hp-Hb complex is formed, the iron becomes unavailable for

bacteria which explains the bacteriostatic effect of Hp (Ceciliani *et al.*, 2012).

It is important to note that Hp may inhibit mast cell proliferation and directly interact the effector cells by binding to CD11/CD18 receptor and also to suppress T-cell proliferation (by inhibition of Th2 response) which expresses its immunomodulatory and anti-inflammatory role (El Ghmati *et al.*, 1996; Murata *et al.*, 2004).

#### *Serum amyloid A*

Serum amyloid A (SAA) is also a major APP in small ruminants (Table 2). Its function is not fully understood but some activities have been well described and documented. Several effects of SAA are well explored, e.g. binding, transporting and scavenging of cholesterol from destroyed cells and lipid debris from bacteria to the liver during inflammation (Ceciliani *et al.*, 2012); inhibition of phagocyte oxidative burst (Linke *et al.*, 1991; Gruys *et al.*, 2005), inhibition of platelet aggregation (Petersen *et al.*, 2004), detoxification of endotoxin (binding to lipopolysaccharide) (Schroedl *et al.*, 2001; Murata *et al.*, 2004), binding to Gram negative bacteria and opsonisation (Hari-Dass *et al.*, 2005; Tothova *et al.*, 2011), inhibition of lymphocytes and endothelial cells proliferation (Murata *et al.*, 2004). It exhibits chemotactic recruitment of inflammatory cells to localised areas of inflammation (Xu *et al.*, 1995; Uhlar & Whitehead, 1999) and down regulates some systemic (pyrexia) events during AP (Shaikin-Kestenbaum *et al.*, 1991; Uhlar & Whitehead, 1999) (Table 1). SAA is synthesised by IL-1, IL-6 and TNF $\alpha$  mainly from hepatocytes but intestinal epithelial cells, macrophages and smooth-muscle cells can also release it by the same pro-inflammatory cytokines (Vreugdenhil *et al.*, 1999; McDonald *et al.*,

2001; Murata *et al.*, 2004). Based on the abovementioned information, it can be concluded that SAA plays a role as an opsonin, prevents accumulation of cholesterol on the site of inflammation and modulates innate immune reactions.

It is established that the normal ovine colostrum contains an isoform of SAA (mammary-associated SAA) which stimulates mucin secretion from the intestine and prevents bacteria development and adapts postparturient period of the newborn animals (McDonald *et al.*, 2001; Mack *et al.*, 2003).

#### *Ceruloplasmin*

Ceruloplasmin (Cp) is a metal-binding APP that stores and transports copper in the body (O'Reilly & Eckersall, 2014) and also plays a role in iron metabolism (Lovstad, 2006) (Table 1). It is classified as a minor APP in sheep and goats (Table 2). Cp is a ferroxidase and acts as an antioxidant that converts the toxic ferrous ion ( $Fe^{2+}$ ) into the non-toxic ferric ion ( $Fe^{3+}$ ) and thus protects the tissues from damaging effects of free radicals (Patel *et al.*, 2002; Murata *et al.*, 2004). Summarised data (Murata *et al.*, 2004) show that Cp is synthesised primarily in the liver but other tissues can be involved including respiratory epithelium of the lung.

#### *Fibrinogen*

Fibrinogen (Fb) is a soluble glycoprotein belonging to  $\beta$ -globulin fraction of the blood plasma. It is classified as a minor APP in sheep but moderate in goats (Table 2). The tissue injury is one of the primary stimulus responsible for activation of the coagulation cascade involving many and different types substances. Fb is a coagulation protein and its synthesis increased during the APR (Davalos & Akassoglou, 2012). It is important to note

that if intravascular coagulation occurs, Fb plasma concentration decreases. Fb is a substrate for fibrin formation and also serves as a matrix for migration of inflammation-related cells and healing of wounds (Raynes, 1994; Thomas, 2000; Murata *et al.*, 2004). It binds specifically to the CD11/CD18 receptors on the migrated phagocytes surface and releases a cascade of intracellular signals that lead to increased degranulation, phagocytosis, antibody-dependent cellular cytotoxicity and delayed apoptosis (Rubel *et al.*, 2001; Murata *et al.*, 2004).

#### *Alfa-1 acid glycoprotein*

Alpha-1 acid glycoprotein (AGP) or orosomucoid is highly glycosylated protein synthesised mainly by liver but extrahepatic production (notably epithelial and endothelial cells) have also been confirmed (Fournier *et al.*, 2000; Murata *et al.*, 2004). AGP is considered as a moderate APP in small ruminants (Table 2). It has at least three important biological functions (Table 1). Like serum albumin, AGP is a binding protein in plasma (O'Reilly & Eckersall, 2014). In physiological conditions, AGP is able to bind more than 300 different biologically active endogenous and exogenous substances such as heparin, histamine, serotonin, steroids, catecholamines and drugs (Fournier *et al.*, 2000; Israili & Dayton, 2001; Eckersall, 2008; Ceciliani *et al.*, 2012). Other suggested AGP functions are the inhibition of neutrophil activation, phagocytosis, platelet activating factor, natural killer cell activity and also a role in T- and B-cell maturation which determines its anti-inflammatory and immunomodulatory activities (Okumura *et al.*, 1985; Fournier *et al.*, 2000; Israili & Dayton, 2001; Murata *et al.*, 2004; Eckersall, 2008). It exhibits a moderate response in

most animals including small ruminants and is likely to be associated with chronic conditions (Tothova *et al.*, 2014). AGP has a role in the innate defense and may act as a non-specific antimicrobial agent by binding directly with lipopolysaccharide and neutralising its toxicity (Moore *et al.*, 1997; Murata *et al.*, 2004; Raich, 2012).

#### *Hemopexin*

Hemopexin (Hpx) is a heme-binding plasma glycoprotein consisting of a single polypeptide chain which is synthesised by hepatocytes in response to IL-6 stimulation (Immenschuh *et al.*, 1995). Heme (iron-protoporphyrin IX) is a component of haemoglobin, myoglobin and some enzymes such as cytochrome, heme peroxidase and others (Smith, 1999). It is released mainly after the destruction of red blood cells. The main function of Hpx is to bind free heme and to protect from oxidative stress, therefore, it acts as an antioxidant (Gutteridge, 1995). Together with Hp and transferrin, Hpx plays a role in the iron homeostasis (Delanghe & Langlois, 2001). Other important role of Hpx is to bind nitric oxide (Shipulina *et al.*, 1998) and carbon monoxide (Shaklai *et al.*, 1981) and thus to neutralise their toxicity. Information concerning Hpx in small ruminants is limited but there are some data about its behaviour during protozoan diseases (Sousa Almeida *et al.*, 2012).

#### *Alfa-1 antitrypsin*

Alpha-1 antitrypsin (AAT) is an APP belonging to serpins with a broad spectrum of antiprotease activity (Table 1). AAT is a serine protease inhibitor which protects tissues from neutrophil proteolytic enzymes (mainly elastase and proteinase) at the site of inflammation (Murata *et al.*, 2004; Tothova *et al.*, 2014).

#### *Lactoferrin*

Lactoferrin (LF) is a part of the innate immunity system and represents one of the first defense reactions against pathogens invading via mucosal tissues (Legnard *et al.*, 2005). LF activities affect the growth and proliferation of infectious agents such as bacteria, protozoan, viruses and fungi (Ward *et al.*, 2002). LF is presented in most mucosal secretions such as colostrum, milk, saliva, small intestine secretion, nasal secretion (Tothova *et al.*, 2014). It is known that neutrophils are the main source of plasma LF (Iyer & Lonnerdal, 1993). In healthy organisms the plasma LF concentration is low but during infection or other tissue injuries it rapidly increases due to inflammatory cells activation (Adlerova *et al.*, 2008). For this reason several authors classify LF as an APP (Kanyshkova *et al.*, 2001). LF plays an important role during infection by binding the iron in stable compound (Adlerova *et al.*, 2008). The complex LF-Fe does not disintegrate even at low pH, notably at the site of inflammation due to metabolic changes occurred by pathogen activities (Valenti & Antonini, 2005). It expresses a bacteriostatic effect of LF that makes the iron unavailable from bacterial growth and proliferation (Tothova *et al.*, 2014).

#### *Albumin*

Albumin (Alb) is the main member of negative APP in ruminants (Table 2). It is the most abundant protein in the blood of animals and humans, representing 35–50% of total protein. Alb is responsible for about 75% of the osmotic pressure of plasma and is the main source of amino acids that may be utilised for synthesis of positive APPs during APR (Ceron *et al.*, 2005; Tothova *et al.*, 2014). Furthermore, the initiation of APR triggers down-regulation of Alb production (Gabay &

Kushner, 1999). Beyond its role as a reservoir for amino acids, Alb also serves as a carrier protein for many organic and inorganic biologically active substances such as thyroxin, estrogen, bilirubin, penicillin, cortisol, free fatty acids, calcium, magnesium, drugs and other which is possibly due to the variety of binding sites on the albumin molecule (Nicholson *et al.*, 2000). The values of plasma Alb may be used as an indicator of nutritional status due to its relatively long half-life (Tothova *et al.*, 2014).

#### *Transferrin*

Transferrin (TF) is a glycoprotein belonging to the family of metal binding transport proteins, synthesised in the liver and together with Alb described as a negative APP (Table 1). TF has a high capability to bind ferric ions in pH of 7.4 but reversibly when the acidity increase (Gomme & McCann, 2005). According to Broch *et al.* (1987), Kaplan *et al.* (1991) and Gkouvatzos *et al.* (2012), iron chelation by transferrin has some important functions such as maintaining Fe<sup>3+</sup> in soluble form under physiological conditions; supporting iron transport and cellular uptake; maintaining Fe<sup>3+</sup> in a redox-inert state, preventing the generation of toxic free radicals and inhibition of the proliferation and growth of bacteria by limiting the access to iron.

#### *C-reactive protein*

C-reactive protein (CPR) is an APP belonging to the pentameric protein family (pentraxins) (Bottazzi *et al.*, 1997). It performs several important functions such as the ability to activate the complement cascade by the classical pathway, modulation of activity of blood platelets, erythrocyte aggregation and opsonisation of necrotic tissues and cellular debris (Vojtic &

Krajnc, 2000). Khan & Khan (2010) also reported that CRP binds with highest affinity to phosphocholine residues in a calcium dependent manner, apoptotic cells, glycans, somatic components of bacteria, fungi and parasites. A number of physiological functions of CPR and its behaviour during various infectious and non-infectious diseases in sheep and goats are not fully elucidated. However, there are data showing that the CRP plasma levels are increased in the case of various types of pneumonia in sheep and goats (Haligur & Ozmen, 2011).

#### *Lipopolysaccharide binding protein*

Lipopolysaccharide binding protein (LBP) is a serum glycoprotein belonging to a small family of lipid-binding proteins together with some other proteins (bactericidal permeability-increasing protein, phospholipid transfer protein, cholesteryl ester transfer protein) (Gustsmann *et al.*, 2001). LBP is synthesised in the liver and intestinal epithelial cells (Ramadori *et al.*, 1990; Vreugdenhil *et al.*, 1999). It plays a key role of innate immune response against bacteria (Ceciliani *et al.*, 2012). Lipopolysaccharide (LPS) molecules are components of external membrane of Gram negative bacteria (Gustsmann *et al.*, 2001) and responsible for the biological toxicity (Raetz, 1990; Rietschel & Brade, 1992). The biological actions of LPS are mediated by both LPS-binding proteins and LPS receptors (Fenton & Golenbock, 1998). LBP binds and transfers a bacterial LPS to the CD14, CD11/18 receptors of antigen-presented cells (Fenton & Golenbock, 1998) and granulocytes (Lamping *et al.*, 1996). This leads to cellular activation resulting in the release of systemically active pro-inflammatory molecules, which in turn mediate systemic toxicity (Fenton & Golenbock, 1998). Enhancement or

inhibition of LPS-induced cellular activation depends on LBP concentration. Low LBP concentration performs a pro-inflammatory role (activation of mononuclear cells), whereas high concentration has an anti-inflammatory role by inhibition LPS-induced cellular stimulation (Lamping *et al.*, 1996; Gustsmann *et al.*, 2001; Ceciliani *et al.*, 2012). LBP is able to bind not only with LPS but also with lipoteichoic acid, a major constituent of the cell wall of Gram-positive bacteria (Ceciliani *et al.*, 2012).

#### ASSESSMENT METHODS AND REFERENCE RANGES

It is known that the plasma levels of APPs are low in the absence of the pathological process. However, there are number of sensitive methods developed for establishing their values both in patients and in

healthy subjects which are presented in Table 3.

Several studies have been conducted in goats and sheep which are designed to determine the reference ranges of APPs in healthy animals (Table 4). In this table are also mentioned some studies that have been conducted in sheep and goats in experimental settings but the data concerning the ranges of APPs values are obtained only from the healthy animals included in the control groups.

#### CONCLUSION

A wide range of investigations have been conducted to determine the usefulness of APPs in various diseases especially in cattle, but data concerning these reactants are still relatively under-utilised in sheep and goats. Overall, they are a useful indi-

**Table 3.** Assays for measuring APPs in small ruminants

APPs	Measurement method	Reference
Hp	Colorimetric method based on the haemoglobin-binding capacity and preservation of the peroxidase activity of the bound haemoglobin at low pH (Phase Haptoglobin Assay kit, Tridelta, Ireland).	Eckersall (2006b)
SAA	Solid phase sandwich ELISA (Tridelta, Ireland)	Colom-Cadena <i>et al.</i> (2014)
Cp	Colorimetric enzyme assay by p-phenylenediamine dihydrochloride oxidation measurement	Meling <i>et al.</i> (2012)
Fb	Heat precipitation method. Nephelometric method.	Ulutas & Ozpinar (2006) Fasulkov <i>et al.</i> (2014)
CPR	Turbidimetric immunoassay by liquid phase immunoprecipitation reaction with rabbit polyclonal antibodies against human CRP (Protiline, BioMérieux)	Vojtic & Krajnc (2000)
ASG	Precipitation method by perchloric acid and determination using bicinchoninic acid protein assay reagent	Eckersall <i>et al.</i> (1996) Gonzalez <i>et al.</i> (2008)
AGP	Radial immunodiffusion by kit (J-Path Inc. Tokyo, Japan).	Eckersall <i>et al.</i> (2007)
LBP	Commercial ELISA kits (Jiancheng Biology Co., Nanjing, China)	Zhong <i>et al.</i> (2014)

**Table 4.** APPs ranges in healthy small ruminants

APPs, species	Normal range (examined)	Reference
Hp, Sheep	0.16±0.13 – 0.19±0.10 g/L	Ulutas & Ozpinar (2006)
	0.197±0.015 g/L	Aziz & Taha (1997)
	0.99±0.12 g/L	Eckersall <i>et al.</i> (2007)
	0.30±0.06 mg/mL	Wells <i>et al.</i> (2013)
	0.038–0.112 g/L	Zhong <i>et al.</i> (2014)
	0–1.0 mg/dL	Jain <i>et al.</i> (2011)
	0.06–0.12 g/L	Lepherd <i>et al.</i> (2009)
	0.048±0.008 mg/mL	Gurdogan <i>et al.</i> (2014)
	0.99 (±0.062) – 0.122 (±0.101) g/L	Nowroozi-Asl <i>et al.</i> (2008)
0.215 (±0.068) – 0.260 (±0.081) mg/mL	Pfeffer & Rogers (1989)	
Hp, Goats	0.12–0.14 mg/mL	Ulutas <i>et al.</i> (2008)
	0.1 g/L	Hashemnia <i>et al.</i> (2011)
	3.05±1.08 g/L	Fasulkov <i>et al.</i> (2014)
	0–0.05 g/L	Gonzalez <i>et al.</i> (2008)
	0.39–1.26 (0.784) mg/dL	Heller & Johns (2015)
	41.6 mg/L	Gonzalez <i>et al.</i> (2011)
	0.58 mg/L	Rahman <i>et al.</i> (2010)
0.056±0.009 mg/L	Balikci <i>et al.</i> (2013)	
SAA, sheep	2.67±0.23 mg/L	Eckersall <i>et al.</i> (2007)
	0.82±0.53 µg/mL	Wells <i>et al.</i> (2013)
	0.963–8.54 mg/L	Zhong <i>et al.</i> (2014)
	0–2.0 µg/mL	Lepherd <i>et al.</i> (2009)
5.62±0.84 µg/mL	Gurdogan <i>et al.</i> (2014)	
SAA, goats	4.20–12.65 µg/mL	Ulutas <i>et al.</i> (2008)
	6.1 µg/mL	Hashemnia <i>et al.</i> (2011)
	1.69–11.94 (4.88) mg/L	Gonzalez <i>et al.</i> (2008)
	0.42–2.2 (0.92) µg/mL	Heller & Johns (2015)
	8.7 µg/mL	Rahman <i>et al.</i> (2010)
6.06±0.89 µg/mL	Balikci <i>et al.</i> (2013)	
AGP, sheep	2.14±0.24 g/L	Eckersall <i>et al.</i> (2007)
	0.246–0.470 g/L	Zhong <i>et al.</i> (2014)
AGP, goats	4.6–40.5 (18.43) mg/dL	Heller & Johns (2015)
LBP, sheep	1.20–2.79 g/L	Zhong <i>et al.</i> (2014)
LBP, goats	0–0.68 (0.125) ng/mL	Heller Johns (2015)
Cp, sheep	11.8±2.3 – 13.5±1.8 mg/dL	Ulutas & Ozpinar (2006)
	0.030–0.039 g/dL	Sousa Almeida <i>et al.</i> (2012)
Cp, goats	4.97–11.85 mg/dL	Ulutas <i>et al.</i> (2008)
	1.1 mg/L	Rahman <i>et al.</i> (2010)
CRP, sheep	0.51±0.19 – 0.55±0.10 mg/dL	Ulutas & Ozpinar (2006)
	5.235±0.429 – 9.115±0.647 mg/L	Vojtic & Krajnc (2000)
Fb, sheep	436.2±39.5 – 437.5±47.8 mg/dL	Ulutas & Ozpinar (2006)
	1.6–4.1 g/L	Lepherd <i>et al.</i> (2009)
Fb, goats	3.2±0.69 g/L	Fasulkov <i>et al.</i> (2014)
	2–4 (2) g/L	Gonzalez <i>et al.</i> (2008)
	2.67 g/L	Gonzalez <i>et al.</i> (2011)
Hpx, sheep	0.013–0.018 g/dl	Sousa Almeida <i>et al.</i> (2012)
ASG, goats	0.86–1.78 (1.25) g/L	Gonzalez <i>et al.</i> (2008)
	1.38 g/L	Gonzalez <i>et al.</i> (2011)
TF, sheep	0.352–0.448 g/dL	Sousa Almeida <i>et al.</i> (2012)
AAT, sheep	0.155–0.193 g/dL	Sousa Almeida <i>et al.</i> (2012)

cator for detecting animals with subclinical infections, determining the prognosis of clinical infection, differentiation between viral and bacterial disease, treatment monitoring, vaccine effectiveness and stress conditions. It is known that the increased APPs concentration is not specific for a particular disease but reflects the overall health of the animals. Based on these data it can be concluded that the knowledge of APPs could be useful for establishment of herd health in small ruminants.

## REFERENCES

- Adlerova, L., A. Bartoskova & M. Faldyna, 2008. Lactoferrin: A review. *Veterinarni Medicina*, **53**, 457–468.
- Aziz, D. M. & M. B. Taha, 1997. Effect of dystocia on serum haptoglobin in Awassi ewes. *Theriogenology*, **48**, 559–562.
- Balikci, E., A. Yildiz & F. Gurdogan, 2013. Selected acute phase proteins, oxidative stress biomarkers, and antioxidants in aborting and non-aborting goats infected with Border disease virus. *Bulletin of the Veterinary Institute in Pulawy*, **57**, 371–374.
- Baumann, H. & J. Gauldie, 1994. The acute phase response. *Immunology Today*, **15**, 74–80.
- Bottazzi, B., V. Vouret-Craviari, A. Bastone, L. De Gioia, C. Matteucci, G. Peri, F. Spreafico, M. Pausa, C. D'Ettorre, E. Giannazza, A. Tagliabue, M. Salmona, F. Tedesco, M. Introna & A. Mantovani, 1997. Multimer formation and ligand recognition by the long pentraxin PTX3. Similarities and differences with the short pentraxins C-reactive protein and serum amyloid P component. *Journal of Biological Chemistry*, **272**, 32817–32823.
- Brock, J. H., T. Mainou-Fowler & S. J. McGregor, 1987. Transferrins and defence against infection. *Annali dell'Istituto Superiore di Sanita*, **23**, 935–941.
- Ceciliani, F., A. Giordano & V. Spagnolo, 2002. The systemic reaction during inflammation: the acute-phase proteins. *Protein and Peptide Letters*, **9**, 211–223.
- Ceciliani, F., J. J. Ceron P. D. Eckersall & H. Sauerwein, 2012. Acute phase proteins in ruminants. *Journal of Proteomics*, **75**, 4207–4231.
- Ceron, J. J., P. D. Eckersall & A. Martinez-Subiela, 2005. Acute phase proteins in dogs and cats: Current knowledge and future perspectives. *Veterinary Clinical Pathology*, **34**, 85–99.
- Colom-Cadena, A., R. Velarde, J. Salinas, K. Borge, I. Garcia-Bocanegra, E. Serrano, D. Gasso, E. Bach, E. Casas-Diaz, J. Lopez-Olivera, S. Lavin, L. Len-Vizcaino & G. Mentaberre, 2014. Management of a caseous lymphadenitis outbreak in a new Iberian ibex (*Capra pyrenaica*) stock reservoir. *Acta Veterinaria Scandinavica*, **56**, 1–11.
- Cray, C., J. Zaias & N. H. Altman, 2009. Acute phase response in animals: A review. *Comparative Medicine*, **59**, 517–526.
- D'Armiento, J., S. Dalal & K. Shada, 1997. Tissue, temporal and inducible expression pattern of haptoglobin in mice. *Gene*, **195**, 19–27.
- Davalos, D. & K. Akassoglou, 2012. Fibrinogen as a key regulator of inflammation in disease. *Seminars in Immunopathology*, **34**, 43–62.
- Delanghe, J. & M. Langlois, 2001. Hemopexin: A review of biological aspects and the role in laboratory medicine. *Clinica Chimica Acta*, **312**, 13–23.
- Eckersall, P. D., P. K. Saini & C. McComb, 1996. The acute phase response of acid soluble glycoprotein,  $\alpha_1$ -acid glycoprotein, ceruloplasmin, haptoglobin and C-reactive protein, in the pig. *Veterinary Immunology and Immunopathology*, **51**, 377–385.
- Eckersall, P. D., 2006a. Measurement of acute phase proteins as biomarkers of disease in production animals. In: *Proceedings of the 57<sup>th</sup> Annual Meeting of the American College of Veterinary Pathologists and the*

- 41<sup>st</sup> Annual Meeting of the American Society for Veterinary Clinical Pathology, Tucson, Arizona, USA.
- Eckersall, P. D., 2006b. Acute phase proteins: form, function and analysis. In: *Proceedings of the 57<sup>th</sup> Annual Meeting of the American College of Veterinary Pathologists and the 41<sup>st</sup> Annual Meeting of the American Society for Veterinary Clinical Pathology*, Tucson, Arizona, USA.
- Eckersall, P. D., 2006c. Acute phase proteins as biomarkers of disease in companion and laboratory animals. In: *Proceedings of the 57<sup>th</sup> Annual Meeting of the American College of Veterinary Pathologists and the 41<sup>st</sup> Annual Meeting of the American Society for Veterinary Clinical Pathology*, Tucson, Arizona, pp 47-52.
- Eckersall, P., F. Lawson, L. Bence, M. Waterston, T. Lang, W. Donachie & M. Fontaine, 2007. Acute phase protein response in an experimental model of ovine caseous lymphadenitis. *BMC Veterinary Research*, **3**, 1–6.
- Eckersall, P. D., 2008. Proteins, proteomics, and the dysproteinemia. In: *Clinical Biochemistry and Domestic animals*, 6<sup>th</sup> edn, eds Kaneko, J. J., J. W. Harvey & M. L. Briss, San Diego, California, USA, pp. 117–155.
- Eckersall, P. D. & R. Bell, 2010. Acute phase proteins: Biomarkers of infection and inflammation in veterinary medicine. *The Veterinary Journal*, **185**, 23–27.
- El Ghmati, S. M., E. M. van Hoeyveld, J. G van Strijp, J. L. Ceuppens & E. A. Stevens, 1996. Identification of haptoglobin as an alternative ligand for CD11B/CD18. *Journal of Immunology*, **156**, 2542–2552.
- Fasulkov, I., M. Karadaev, N. Vasilev, V. Urumova & T. Mircheva, 2014. Determination of plasma fibrinogen and haptoglobin, hematological and blood biochemical changes in Bulgarian local goats with experimentally induced *Staphylococcus aureus* mastitis. *Turkish Journal of Veterinary and Animal Sciences* **38**, 439–444.
- Fenton, M. & D. Golenbock, 1998. LPS-binding proteins and receptors. *Journal of Leukocyte Biology*, **64**, 25–32.
- Fournier, T., N. Medjoubi-N & D. Porquet, 2000. Alpha-1-acid glycoprotein. *Biochimica et Biophysica Acta*, **1482**, 157–171.
- Gabay, C. & I. Kushner, 1999. Acute-phase proteins and other systemic responses to inflammation. *New England Journal of Medicine*, **340**, 448–454.
- Gkouvatsos, K., G. Papanikolaou & K. Pantopoulos, 2012. Regulation of iron transport and the role of transferrin. *Biochimica et Biophysica Acta*, **1820**, 188–202.
- Gomme, P. T. & K. B. McCann, 2005. Transferrin: Structure, functional and potential therapeutic action. *Drug Discovery Today*, **10**, 267–273.
- Gomez-Laguna, J. F. Salguero, F. Pallares, I. Rodriguez-Gómez, I. Barranco & L. Carrasco, 2011. Acute phase proteins as biomarkers in animal health and welfare. In: *Acute Phase Proteins as Early Non-Specific Biomarkers of Human and Veterinary Diseases*, ed F. Veas, InTech, pp. 259–298.
- Gonzalez, F. H., F. Tecles, S. Martinez-Subiela, A. Tvarijonaviute, L. Soler & J. J. Ceron, 2008. Acute phase protein response in goats. *Journal of Veterinary Diagnostic Investigation*, **20**, 580–584.
- Gonzalez, F., F. Hernandez, J. Madrid, S. Martinez-Subiela, A. Tvarijonaviute, J. Ceron & F. Tecles, 2011. Acute phase proteins in experimentally induced pregnancy toxemia in goats. *Journal of Veterinary Diagnostic Investigation*, **23**, 57–62.
- Gruys, E., M. J. M. Toussaint, W. J. M. Landman, M. Tivapasi, R. Chamanza & L. van Veen, 1999. Infection, inflammation and stress inhibit growth. Mechanisms and non-specific assessment of the processes by acute phase proteins. In: *Production Diseases in Farm Animals: 10<sup>th</sup> International Conference*, ed T. Wensing, Wageningen Press, Wageningen, pp. 72–87.
- Gruys, E., M. J. M. Toussaint, T. A. Niewold & S. J. Koopmans, 2005. Acute phase re-

- action and acute phase proteins. *Journal of Zhejiang University Science B*, **6**, 1045–1056.
- Gurdogan, F., E. Balıkcı & A. Yildiz, 2014. Some acute phase proteins, oxidative stress biomarkers and antioxidant enzyme activities in ewes with pregnancy toxemia. *Iranian Journal of Veterinary Research*, **15**, 297–299.
- Gutteridge J. M. C., 1995. Lipid peroxidation and antioxidants as biomarkers of tissue damage. *Clinical Chemistry*, **41**, 1819–1828.
- Gutsmann, T., M. Müller, R. MacKenzie, A. Wiese & U. Seydel, 2001. Dual role of lipopolysaccharide (LPS)-binding protein in neutralization of LPS and enhancement of LPS-induced activation of mononuclear cells. *Infection and Immunity*, **69**, 6942–6950.
- Haligur, M. & O. Özmen, 2011. Immunohistochemical detection of serum amyloid-A, serum amyloid-P, C-reactive protein, tumour necrosis factor- $\alpha$  and TNF- $\alpha$  receptor in sheep and goat pneumonias. *Révue de Médecine Vétérinaire*, **162**, 475–481.
- Hari-Dass, R., C. Shah, D. J. Meyer & J. G. Raynes, 2005. Serum amyloid A protein binds to outer membrane protein A of Gram-negative bacteria. *Journal of Biological Chemistry*, **280**, 18562–18567.
- Hashemnia, M., A. Khodakaram-Tafti, S. Razavi & S. Nazifi, 2011. Changing patterns of acute phase proteins and inflammatory mediators in experimental caprine coccidiosis. *Korean Journal of Parasitology*, **49**, 213–219.
- Heegaard, P. M., D. L. Godson, M. J. Toussaint, K. Tjønehoj, L. E. Larsen, B. Viuff & L. Ronsholt, 2000. The acute phase response of haptoglobin and serum amyloid A (SAA) in cattle undergoing experimental infection with bovine respiratory syncytial virus. *Veterinary Immunology and Immunopathology*, **77**, 9–15.
- Heller, M. & J. Johns, 2015. Acute phase proteins in healthy goats: Establishment of reference intervals. *Journal of Veterinary Diagnostic Investigation*, **27**, 171–181.
- Immenschuh, S., D. X. Song, H. Satoh & U. Müller-Eberhard, 1995. The type II hemopexin interleukin-6 response element predominates the transcriptional regulation of the hemopexin acute phase responsiveness. *Biochemical and Biophysical Research Communications*, **207**, 202–208.
- Israili, Z. H. & P. G. Dayton, 2001. Human alpha-1-glycoprotein and its interactions with drugs. *Drug Metabolites Review*, **33**, 161–235.
- Iyer, S. & B. Lonnerdal, 1993. Lactoferrin, lactoferrin receptors and iron metabolism. *European Journal of Clinical Nutrition*, **47**, 232–241.
- Jain, S., V. Gautam & S. Naseem, 2011. Acute-phase proteins: A diagnostic tool. *Journal of Pharmacy and Bioallied Science*, **3**, 118–127.
- Janeway, C. A., P. Travers, M. Walport & M. J. Schlomschik, 2001. Immunology, 5<sup>th</sup> edn, Taylor and Francis, London, pp. 732.
- Kanyshkova, T. G., V. N. Buneva & G. A. Nevinsky, 2001. Lactoferrin and its biological functions. *Biochemistry (Moscow)*, **66**, 1–7 (RU).
- Kaplan, J., I. Jordan & A. Sturrock, 1991. Regulation of the transferrin-independent iron transport system in cultured cells. *Journal of Biological Chemistry*, **266**, 2997–3004.
- Kato, G., 2009. Haptoglobin halts haemoglobin's havoc. *Journal of Clinical Investigation*, **119**, 2140–2142.
- Khan, F. & M. Khan, 2010. Inflammation and acute phase response. *International Journal of Applied Biology and Pharmaceutical Technology*, **1**, 312–321.
- Lamping, N., A. Hoess, B. Yu, T. Park, S. Kirschning, D. Pfeil, D. Reuter, S. Wright, F. Herrmann & R. Schumann, 1996. Effects of site-directed mutagenesis of basic residues (Arg 94, Lys 95, Lys 99) of lipopolysaccharide (LPS)-binding protein on binding and transfer of LPS and subsequent immune cell activation. *Journal of Immunology*, **157**, 4648–4656.

- Legnard, D., E. Ellass, M. Carpentier & J. Mazurier, 2005. Lactoferrin: A modulator of immune and inflammatory response. *Cellular and Molecular Life Sciences*, **62**, 2549–2559.
- Lepherd, M. L., P. J. Canfield, G. B. Hunt & K. L. Bosward, 2009. Haematological, biochemical and selected acute phase protein reference intervals for weaned female Merino lambs. *Australian Veterinary Journal*, **87**, 5–11.
- Linke, R. P., V. Bock, G. Valet & G. Rothe, 1991. Inhibition of the oxidative burst response of N-formyl peptide-stimulated neutrophils by serum amyloid-A protein. *Biochemical and Biophysical Research Communication*, **176**, 1100–1105.
- Lovstad, R. A., 2006. A kinetic study on the phenothiazine dependent oxidation of NADH by bovine ceruloplasmin. *Bio-metals*, **19**, 1–5.
- Mack, D. R., T. L. McDonald, M. A. Larson, S. Wei & A. Weber, 2003. The conserved TFLK motif of mammary-associated serum amyloid A is responsible for up-regulation of intestinal MUC3 mucin expression in vitro. *Pediatric Research*, **53**, 137–142.
- McDonald, T. L., M. A. Larson, D. R. Mack & A. Weber, 2001. Elevated extrahepatic expression and secretion of mammary-associated serum amyloid A 3 (M-SAA3) into colostrum. *Veterinary Immunology and Immunopathology*, **83**, 203–211.
- Meling, S., K. Bardsen & M. Ulvund, 2012. Presence of an acute phase response in sheep with clinical classical scrapie. *BMC Veterinary Research*, **8**, 1–11.
- Moore, D. F., M. R. Rosenfeld, P. M. Gribbon, C. P. Winlove & C. M. Tsai, 1997. Alpha-1-acid (AAG, orosomucoid) glycoprotein: Interaction with bacterial lipopolysaccharide and protection from sepsis. *Inflammation*, **21**, 69–82.
- Moshage, H., 1997. Cytokines and the hepatic acute phase response. *Journal of Pathology*, **181**, 257–266.
- Murata, H., N. Shimada & M. Yoshioka, 2004. Current research on acute phase proteins in veterinary diagnosis: an overview. *The Veterinary Journal*, **168**, 28–40.
- Murtaugh, M. P., 1994. Porcine cytokines. *Veterinary Immunology and Immunopathology*, **43**, 37–44.
- Nicholson, J. P., M. R. Wolmarans & G. R. Park, 2000. The role of albumin in critical illness. *British Journal of Anaesthesia*, **85**, 599–610.
- Niewold, T. A., M. J. M. Toussaint & E. Gruys, 2003. Monitoring health by acute phase proteins. In: *Proceedings of the Fourth European Colloquium on Acute Phase Proteins*, Segovia, Spain, pp. 57–67.
- Nowroozi-Asl, A., S. Nazifi & A. Bahari, 2008. Determination of serum haptoglobin reference value in clinically healthy Iranian fat-tailed sheep. *Iranian Journal of Veterinary Research*, **9**, 171–173.
- Okumura, Y., J. Kudo, T. Ikuta, S. Kurokawa, H. Ishibashi & H. Okubo, 1985. Influence of acute-phase proteins on the activity of natural killer cells. *Inflammation*, **9**, 211–219.
- O'Reilly, E. L. & P. D. Eckersall, 2014. Acute phase proteins: a review of their function, behavior and measurement in chickens. *World's Poultry Science Journal*, **70**, 27–43.
- Pannen, B. H. J. & J. L. Robotham, 1995. The acute-phase response. *New Horizons*, **3**, 183–197.
- Patel, B. N., R. J. Dunn, S. Y. Jeong, Q. Zhu, J. P. Julien & S. David, 2002. Ceruloplasmin regulates iron levels in the CNS and prevents free radical injury. *Journal of Neuroscience*, **22**, 6578–6586.
- Petersen, H. H., J. P., Nielsen & P. M. H. Heegaard, 2004. Application of acute phase protein measurements in veterinary clinical chemistry. *Veterinary Research*, **35**, 163–187.
- Pfeffer, A. & K. M. Rogers, 1989. Acute phase response of sheep: Changes in the concentrations of ceruloplasmin, fibrinogen, haptoglobin and the major blood cell types as

- sociated with pulmonary damage. *Research in Veterinary Science*, **46**, 118–124.
- Pyorala, S., 2000. Hirvonen's thesis on acute phase response in dairy cattle. PhD thesis. University of Helsinki, Helsinki, Finland.
- Raetz, C. R. H., 1990. Biochemistry of endotoxins. *Annual Review of Biochemistry*, **59**, 129–170.
- Rahman, M. M., C. Lecchi, C. Fraquelli, P. Sartorelli & F. Ceciliani, 2010. Acute phase protein response in Alpine ibex with sarcoptic mange. *Veterinary Parasitology*, **168**, 293–298.
- Raich, E., 2012. Acute phase proteins in wild boar, Pyrenean chamois and ibex: Method validation and reference values. *Tesi Doctoral*, Universitat Autònoma de Barcelona, pp. 29–32.
- Ramadori, G., K. Meyer zum Buschenfelde, P. Tobias, J. Mathison & R. Ulevitch, 1990. Biosynthesis of lipopolysaccharide-binding protein in rabbit hepatocytes. *Pathobiology*, **58**, 89–94.
- Raynes, J. D., 1994. The acute phase response. *Biochemical Society Transactions*, **22**, 69–74.
- Rietschel, E. T. & H. Brade, 1992. Bacterial endotoxins. *Scientific American*, **267**, 26–31.
- Rubel, C., G. C. Fernandez, G. Dran, M. B. Bompadre, M. A. Isturiz & M. S. Palermo, 2001. Fibrinogen promotes neutrophil activation and delays apoptosis. *Journal of Immunology*, **166**, 2002–2010.
- Schaer, D. J., F. S. Roberti, G. Schoedon & A. Schaffner, 2002. Induction of the CD163-dependent haemoglobin uptake by macrophages as a novel anti-inflammatory action of glucocorticoids. *British Journal of Haematology*, **119**, 239–243.
- Schroedl, W., B. Fuerll, P. Reinhold, M. Krueger & C. Schuett, 2001. A novel acute phase marker in cattle: lipopolysaccharide binding protein (LBP). *Journal of Endotoxin Research*, **7**, 49–52.
- Shainkin-Kestenbaum, R., G. Berlyne, S. Zimlichman, H. R. Sorin, M. Nyska & A. Dannon, 1991. Acute phase protein, serum amyloid A, inhibits IL-1 and TNF induced fever and hypothalamic PGE<sub>2</sub> in mice. *Scandinavian Journal of Immunology*, **34**, 179–183.
- Shaklai, N., V. S. Sharma, U. Muller-Eberhard & W. T. Morgan, 1981. The interaction of heme-hemopexin with CO. *Journal of Biological Chemistry*, **256**, 1544–1548.
- Shipulina, N., R. C. Hunt & N. Shaklai, A. Smith, 1998. Coordination of nitric oxide by heme-hemopexin. *Journal of Protein Chemistry*, **17**, 255–260.
- Sipe, J. D., 1985. Acute phase proteins in chronic inflammation. In: *The Acute Phase Response to Injury and Infection*. ed A. Koj, Elsevier, Amsterdam, pp. 273–284.
- Smith, D. J. & D. Roberts, 1994. Effects of high volume and/or intense exercise on selected blood chemistry parameters. *Clinical Biochemistry*, **27**, 435–440.
- Smith A., 1999. Role of redox-reactive metals in the regulation of the metallothionein and heme oxygenase genes by heme and hemopexin. In: *Iron Metabolism*, eds G. C. Ferreira, J. J. G. Moura & R. Franco, Wiley-VCH, Weinheim, pp. 65–92.
- Sousa Almeida, K., A. Costa, P. Silva, J. Fagliari, R. Machado & A. Nascimento, 2012. Acute phase proteins: A potential approach for diagnosing chronic infection by *Trypanosoma vivax*. *Revista Brasileira de Parasitologia Veterinaria*, **21**, 97–100.
- Thomas, J. S., 2000. Overview of plasma proteins. In: *Schalm's Veterinary Hematology*, 5<sup>th</sup> edn, eds B. F. Feldman, J. G. Zinkl & N. C. Jain, Lippincott Williams, Wilkins, Philadelphia, pp. 891–898.
- Tirziu, E., 2009. Acute-phase proteins in immune response. *Lucraris Stiintifice Medicina Veterinara*, **42**, 329–339.
- Tothova, C., O. Nagy, H. Seidel & G. Kováč, 2011. Acute phase proteins as markers of diseases in farm animals. In: *Acute Phase Proteins as Early Non-Specific Biomarkers of Human and Veterinary Diseases*, ed F. Veas, InTech, pp. 231–258.
- Tothova, S., O. Nagy & G. Kovac, 2014. Acute phase proteins and their use in the diag-

- nosis of diseases in ruminants: a review. *Veterinarni Medicina*, **59**, 163–180.
- Uhlar, C. M. & A. S. Whitehead, 1999. Serum amyloid A, the major vertebrate acute-phase reactant. *European Journal of Biochemistry*, **265**, 501–523.
- Ulutas, P. & A. Ozpinar, 2006. Effect of *Mannheimia (Pasteurella) haemolytica* infection on acute-phase proteins and some mineral levels in colostrum–breast milk-fed or colostrum–breast milk-deprived sheep. *Veterinary Research Communications*, **30**, 485–495.
- Ulutas, P., H. Voyvoda, B. Ulutas & S. Aypak, 2008. Haptoglobin, serum amyloid- $\alpha$  and ceruloplasmin concentrations in goats with mixed helminth infection. *Türkiye Parazitoloji Dergisi*, **32**, 229–233.
- Ward, P. P., S. Uribe-Luna & O. M. Conneely, 2002. Lactoferrin and host defense. *Biochemistry and Cell Biology*, **80**, 95–102.
- Valenti, P. & G. Antonini, 2005. Lactoferrin: An important host defense against microbial and viral attack. *Cellular and Molecular Life Sciences*, **62**, 2576–2587.
- van Miert, A. S., 1995. Pro-inflammatory cytokines in a ruminant model: pathophysiological, pharmacological, and therapeutic aspects. *Veterinary Quarterly* **175**, 41–50.
- Vojtic, I. & S. Krajnc, 2000. Determination of C-reactive protein by turbidimetric immunoassay (TIA) in sheep. *Veterinarski Arhiv*, **70**, 151–157.
- Vreugdenhil, A. C., M. A. Dentener, A. M. Snoek, J. W. Greve & W. A. Buurman, 1999. Lipopolysaccharide binding protein and serum amyloid A secretion by human intestinal epithelial cells during the acute phase response. *Journal of Immunology*, **163**, 2792–2798.
- Wagener, F. A., A. Eggert, O. C. Boerman, W. J. Oyen, A. Verhofstad, N. G. Abraham, G. Adema, Y. Van Kooyk, T. De Witte & C. G. Figdor, 2001. Heme is a potent inducer of inflammation in mice and is counteracted by heme oxygenase. *Blood*, **98**, 1802–1811.
- Wells, B., I. Innocent, P. Eckersall, E. McCulloch, A. Nisbet & S. Burgess, 2013. Two major ruminant acute phase proteins, haptoglobin and serum amyloid A, as serum biomarkers during active sheep scab infestation. *Veterinary Research*, **44**, 103.
- Xu, L., R. Badolato, W. J. Murphy, D. L. Longo, M. Anver, S. Hale, J. J. Oppenheim & J. M. Wang, 1995. A novel biologic function of serum amyloid A. Induction of T lymphocyte migration and adhesion. *Journal of Immunology*, **155**, 1184–1190.
- Yang, F., D. J. Haile, F. G. Berger, D. C. Herbert, E. Van Beveren & A. J. Ghio, 2003. Haptoglobin reduces lung injury associated with exposure to blood. *American Journal of Physiology – Lung and Cell Molecular Physiology*, **284**, 402–409.
- Zhong, R., H. Yang Li, H. Sun & D. Zhou, 2014. Effects of supplementation with dietary green tea polyphenols on parasite resistance and acute phase protein response to *Haemonchus contortus* infection in lambs. *Veterinary Parasitology*, **205**, 199–207.

Paper received 09.06.2016; accepted for publication 30.09.2016

**Correspondence:**

Petar T. Iliev  
Department of Veterinary Microbiology,  
Infectious and Parasitic Diseases,  
Faculty of Veterinary Medicine,  
Trakia University  
tel: +359 42 699 587  
e-mail: petyo\_todorow@abv.bg