



# Regional effects of atmospheric aerosols on temperature: an evaluation of an ensemble of on-line coupled models

**Rocío Baró<sup>1</sup>, Laura Palacios-Peña<sup>1</sup>, Alexander Baklanov<sup>2</sup>, Alessandra Balzarini<sup>3</sup>, Dominik Brunner<sup>4</sup>, Renate Forkel<sup>5</sup>, Marcus Hirtl<sup>6</sup>, Luka Honzak<sup>7</sup>, Juan Luis Pérez<sup>8</sup>, Guido Pirovano<sup>3</sup>, Roberto San José<sup>8</sup>, Wolfram Schröder<sup>9</sup>, Johannes Werhahn<sup>5</sup>, Ralf Wolke<sup>9</sup>, Rahela Zabkar<sup>10</sup>, and Pedro Jiménez-Guerrero<sup>1</sup>**

<sup>1</sup>Department of Physics, Regional Campus of International Excellence Campus Mare Nostrum, University of Murcia, Murcia, Spain

<sup>2</sup>World Meteorological Organization, Geneva, Switzerland

<sup>3</sup>Ricerca sul Sistema Energetico (RSE), Italy

<sup>4</sup>Laboratory for Air Pollution/Environmental Technology, Empa, Swiss Federal Laboratories for Materials Science and Technology, Switzerland

<sup>5</sup>Karlsruher Institut für Technologie (KIT), Institut für Meteorologie und Klimaforschung, Atmosphärische Umweltforschung (IMK-IFU), Germany

<sup>6</sup>Section Chemical Weather Forecasts, Division Data/Methods/Modeling, ZAMG - Zentralanstalt für Meteorologie und Geodynamik, Austria

<sup>7</sup>BO-MO d.o.o, Slovenia

<sup>8</sup>Environmental Software and Modelling Group, Computer Science School - Technical University of Madrid, Spain

<sup>9</sup>Leibniz Institute for Tropospheric Research, Permoserstr, Germany

<sup>10</sup>Slovenian Environment Agency, Slovenia

Correspondence to: Pedro Jiménez-Guerrero (pedro.jimenezguerrero@um.es)



## Abstract

The climate effect of atmospheric aerosols is associated to their influence on the radiative budget of the Earth due to direct aerosol-radiation interactions (ARI) and indirect effects, resulting from aerosol-cloud interactions (ACI). On-line coupled meteorology-chemistry models permit the description of these effects on the basis of simulated atmospheric aerosol concentrations, although there is still some uncertainty associated to the use of these models. In this sense, the objective of this work is to assess whether the inclusion of atmospheric aerosol radiative feedbacks of an ensemble of on-line coupled models improves the simulation results for maximum, mean and minimum temperature over Europe. The evaluated model outputs originate from EuMetChem COST Action ES1004 simulations for Europe, differing in the inclusion (or omission) of ARI and ACI in the various models. The case studies cover two important atmospheric aerosol episodes over Europe in the year 2010, a heat wave and forest fires episode (July-August 2010) and a more humid episode including a Saharan desert dust outbreak in October 2010. The simulation results are evaluated against observational data from E-OBS gridded database. The results indicate that, although there is only a slight improvement in the bias of the simulation results when including the radiative feedbacks, the spatio-temporal variability and correlation coefficients are improved for the cases under study when atmospheric aerosol radiative effects are included, especially for those areas closest to emissions sources of atmospheric aerosols.

## 1 Introduction

Atmospheric aerosol particles are known to have an impact on Earth's radiative budget due to their optical, microphysical and chemical properties, and are considered to be the most uncertain forcing agent. They influence climate by modifying both the global energy balance through absorption and scattering of radiation (direct effect), and by acting as cloud condensation nuclei, thus affecting cloud droplet size distributions and lifetime (Twomey, 1977; Lohmann and Feichter, 2005; Chung, 2012) and the reflectance and persistence of



clouds (indirect effects) (Ghan and Schwartz, 2007; Yang et al., 2011). Depending on the atmospheric aerosol concentration, aerosol cloud interactions may result in an increase or decrease in liquid water content, cloud cover, and lifetime of low level clouds and a suppression or enhancement of precipitation (Bangert et al., 2011). Besides, aerosol absorption may decrease low-cloud cover by heating the air and reducing relative humidity. This leads to a positive radiative forcing, termed the semi-direct effect, which amplifies the warming influence of absorbing aerosols (Hansen et al., 1997). The Fifth Report of the Intergovernmental Panel on Climate Change (IPCC AR5) (Boucher et al., 2013; Myhre et al., 2013) distinguishes between aerosol-radiation interactions (ARI), which encompass the aerosol direct and semidirect effect, and the aerosol-cloud interactions (ACI), which encompass the indirect effects.

In order to account for these atmospheric aerosol effects, the use of fully-coupled models is needed for meteorological, chemical and physical processes. On-line coupled models include the interaction of atmospheric pollutants (gaseous-phase compounds and aerosols) with meteorological variables (Baklanov et al., 2014). In this context, the air quality model evaluation international initiative (AQMEII) in its phase 2 (Alapaty et al., 2012; Galmarini et al., 2015) focused on the assessment of how well the current generation of coupled regional scale air quality models can simulate the spatio-temporal variability in the optical and radiative characteristics of atmospheric aerosols and associated feedbacks among aerosols, radiation, clouds, and precipitation. On this basis, a coordinated exercise of Working Groups 2 and 4 of the COST Action ES1004 (EuMetChem, <http://eumetchem.info>) emerged, in order to take into account the radiative feedbacks, due to atmospheric aerosol effects over meteorology. In this initiative, two important episodes with high loads of atmospheric aerosols were analyzed which were identified during the previous AQMEII phase 2 modelling intercomparison exercise (Galmarini et al., 2015). They were selected on behalf of their strong potential of aerosol interactions (Makar et al., 2015a, b; Forkel et al., 2015).

As a result of the AQMEII phase 2 initiative and EuMetChem COST Action, several studies covering the analysis of the ARI+ACI feedbacks onto meteorology have been done (e.g. Baró et al. (2015); Forkel et al. (2015, 2016); Kong et al. (2015); San José et al. (2015)).



Focusing on the effects of including ARI+ACI on temperature, Forkel et al. (2015) focused on the 2010 Russian wildfire episode, where the presence of the atmospheric aerosols decreased the mean temperature during summer 2010 by 0.25 K over the target area. For the same episode, Péré et al. (2014) showed daily mean surface temperature reductions between 0.2 to 2.6 K. Forkel et al. (2012) studied an episode in June and July 2006, where a slightly lower temperature was found over western Europe when including atmospheric aerosol feedbacks; this reduction was reflected in its spatial distribution of the planetary boundary layer height. Moreover, Meier et al. (2012) found during July 2006 a general decrease of 0.14 K on 2-m temperature when simulating absorbing aerosol in upper layers compared to an aerosol-free troposphere over land surface.

However, all these studies are based on individual model evaluations and do not take into account an ensemble of regional models, in order to build confidence on model simulations and to characterize the uncertainty associated to the use of different modelling systems. Therefore, the objective of this work is to assess whether the inclusion of aerosol radiative feedbacks during two important atmospheric aerosol episodes of the year 2010 improves the outputs of an ensemble of regional on-line coupled models for maximum, mean and minimum temperature at 2 meters over Europe.

## 2 Methodology

The analyzed model outputs are the results of a coordinated modeling exercise which was performed within the COST Action ES1004 (EuMetChem). In order to analyze the ARI or ARI+ACI effect on temperature, it was suggested to run three case studies for two episodes are run with different on-line coupled models with identical meteorological boundary conditions and anthropogenic emissions. The two considered episodes are: the Russian 2010 heatwave and wildfires episode in summer 2010 (25 July-15 August 2010) and an autumn Saharan dust episode, including the dust transport to Europe (2-15 October 2010).

For the chosen episodes, simulations with each model were performed with and without considering the atmospheric aerosol effects. Three different cases were requested: the first



one which does not consider any feedbacks to meteorology with simulated aerosol (NRF); second, where only direct aerosol effect is considered (ARI) and third, where aerosol-cloud interactions based on simulated aerosol concentrations and direct and indirect aerosol effects are considered (ARI+ACI). However, the third case could not be submitted by all of the participants. The common setup for the participating models and a unified output strategy allow analyzing the model output with respect to similarities and differences in the model response to the aerosol direct effect and aerosol-cloud interactions.

## 2.1 Participating models

An overview of the different models and their configurations is shown in Table 1. The participating models shown here are COSMO-MUSCAT (Wolke et al., 2012) and WRF-Chem (Grell et al., 2005; Grell and Baklanov, 2011) with different chemistry and physics options and performed episodes. The horizontal resolution is around 25 km for most of the contributions. Only for the fire episode, the COSMO-MUSCAT simulations were made with a grid with of 0,125deg /approximately 14 km) there is an additional WRF-Chem run with 9 km width. The COSMO models use Kessler-type bulk microphysics (Doms et al., 2011) and WRF-Chem uses Morrison microphysics (Morrison et al., 2009), except for one contribution, that utilizes Lin (Lin et al., 1983). COSMO models use prognostic TKE (Doms et al., 2011) planetary boundary layer (PBL). The YSU PBL scheme (Hong et al., 2006) was chosen for the WRF-Chem simulations. In general, the Modal Aerosol Dynamics Model for Europe (MADE) is applied (Ackermann et al., 1998) except for one WRF-Chem simulation, which uses the Model for Simulating Aerosol Interactions and Chemistry (MOSAIC)(4 bins) approach (Zaveri et al., 2008). For further information and details about the models, we refer to the work of Forkel et al. (2015); Im et al. (2015a, b); Baró et al. (2015). To enable the cross-comparison between models, the participating groups interpolated their model output to a common grid with 0.1 degree resolution.

Moreover, the ensemble of the available simulations has also been included in this comparison, as recommended by several studies (Vautard et al., 2012; Jiménez-Guerrero et al., 2013; Landgren et al., 2014; Solazzo and Galmarini, 2015; Kioutsioukis et al., 2016), in or-



der to check whether the design of an ensemble of simulations outperforms (or not) the skill of individual models.

## 2.2 Emissions and boundary conditions

For the EU domain, the anthropogenic emissions for the year 2009 (<http://www.gmes-atmosphere.eu/>) were applied by all modelling groups and are based on the TNO-MACC-II (Netherlands Organization for Applied Scientific Research, Monitoring Atmospheric Composition and Climate–Interim Implementation) framework (Kuenen et al., 2014; Pouliot et al., 2015). As described in Im et al. (2015a), annual emissions of methane (CH<sub>4</sub>), carbon monoxide (CO), ammonia (NH<sub>3</sub>), total non-methane volatile organic compounds (NMVOC), nitrogen oxides (NO<sub>x</sub>), particulate matter (PM<sub>10</sub> & PM<sub>2.5</sub>) and sulfur dioxide (SO<sub>2</sub>) from ten activity sectors are provided on a latitude/longitude grid of 1/8 × 1/16 resolution. For further details, the reader is referred to Im et al. (2015a, b).

Hourly biomass burning emissions were provided by the Finnish Meteorological Institute (FMI) fire assimilation system (<http://is4fires.fmi.fi/>) (Sofiev et al., 2009). More details on the fire emissions and their uncertainties are discussed in Soares et al. (2015). The fire assimilation system provides only data for total PM emissions; the estimation of emissions for other species are described in Im et al. (2015b).

## 2.3 Observational database

E-OBS (Haylock et al., 2008) version 11.0 has been used as the gridded observational database for maximum, mean and minimum temperature. E-OBS is a high-resolution European land-only daily gridded data set covering the period 1950-2014. The E-OBS 0.25 degrees regular latitude-longitude grid has been used as the reference for validation. Thus, data from all model runs have been bilinearly interpolated onto the E-OBS grid. Since the resolution of the models is similar to that of E-OBS, the interpolation procedure is not expected to alter significantly any of our results.



## 2.4 Validation methodology

All the statistical measures are calculated at individual grid points. Only land grid points are considered in the analysis, since these are the only points where E-OBS contains information. Areas in grey indicate cells where E-OBS data are not available (southeastern part of the domain for the wildfires or southern part of the domain in the dust episode) or areas not covered by the modelling domain (southern part of the domain for the CS2 configuration).

We will use the notation  $V_{ipc}^k$  for a variable from model  $k$  at grid point  $i$ , on period  $p$ =fires,dust and case  $c$ =1 2, 3 representing no radiative feedbacks, ARI and ARI+ACI. If we use bracket notation for an average over a given index (e.g.  $\langle \cdot \rangle_{pc}$ ), we can express the bias at a given grid point as:

$$b_i^k = \left\langle V_{ipc}^k - O_{ip} \right\rangle_{pc} \quad (1)$$

where  $O_{ip}$  is the value observed. The model bias is the simplest measure of model performance.

The ensemble mean,  $\left\langle V_{ipc}^k \right\rangle_k$ , is usually considered as an additional simulation which compensates the errors of the different ensemble members. Even though this is a very simplistic view of the ensemble (which should be considered from a probabilistic point of view), it can be useful to reinforce the common signal of the different models in our analysis of the mean climate. Notice, however, that the ensemble mean is not a physical realization of any of the models, but just a statistical average (Knutti et al., 2010; Jiménez-Guerrero et al., 2013).

Then, the variability was assessed on the hourly series ( $\hat{V}_{ipc}^k$ ):

$$\hat{V}_{ipc}^k = V_{ipc}^k - V_{ic}^k \quad (2)$$

The ability to represent the variability can be decomposed into:



- the ability to represent its size, which can be represented by the standard deviation of the series:

$$\text{sd}[V]_i^k = \sqrt{\left\langle \left( \hat{V}_{ipc}^k \right)^2 \right\rangle_{pc}} \quad (3)$$

and can be compared to that of the observations  $\text{sd}[O]_{ip}$ , and

- the ability to represent the hourly variations, which can be represented by the linear determination coefficient ( $\rho^2$ ) with the observations.

$$\rho_i^{2,k} = \frac{\left\langle \hat{V}_{ipc}^k \hat{O}_{ip} \right\rangle_{pm}^2}{\left\langle \left( \hat{V}_{ipc}^k \right)^2 \right\rangle_{pm} \left\langle \left( \hat{O}_{ip} \right)^2 \right\rangle_{pc}} \quad (4)$$

The latter ability can only be expected on simulations nested into “perfect” boundary conditions such as those considered in this study.

Finally, pattern agreement between simulated and observed data was quantified in a Taylor diagram by means of the spatial correlation ( $r$ ) and the ratio between simulated and observed standard deviations,  $V_i^k \equiv \left\langle V_{ipc}^k \right\rangle_{pc}$

$$r^k = \frac{\left\langle \left( V_i^k - \left\langle V_i^k \right\rangle_i \right) \left( O_i - \left\langle O_i \right\rangle_i \right) \right\rangle_i}{\sqrt{\left\langle \left( V_i^k - \left\langle V_i^k \right\rangle_i \right)^2 \right\rangle_i \left\langle \left( O_i - \left\langle O_i \right\rangle_i \right)^2 \right\rangle_i}} \quad (5)$$

$$s^k = \sqrt{\frac{\left\langle \left( V_i^k - \left\langle V_i^k \right\rangle_i \right)^2 \right\rangle_i}{\left\langle \left( O_i - \left\langle O_i \right\rangle_i \right)^2 \right\rangle_i}} \quad (6)$$

This information can be summarized in a Taylor (2001) diagram, which is a polar plot, with radial coordinate  $s^k$  and angular coordinate related to  $r^k$ .



## 3 Results

### 3.1 Bias

The results for the daily bias of maximum, mean and minimum temperature have been obtained by calculating the bias of the daily mean series at each grid point of all the land grid points of the corresponding domain for the fires and dust episodes.

During the fire episode (Fig. 1 left column) there is a general underestimation of the maximum temperature in the base case (average domain values from -2.1 K in ES3-C11 to -1.2 K in DE3-C11). This is especially noticeable over several cells in Russia (up to -7 K). Conversely, a general overestimation is found in the west and northwest area of the domain (positive differences between +1.0 K in DE3-C11 to +6.5 K in ES1-C11). When introducing the ARI or ARI+ACI, model biases do not improve (mean variation of the bias of +17.2% in C12 and +11.0% in C13). This positive variation was expected because the cold bias of models for reproducing maximum temperature and the overall cooling effects of aerosols. However, the improvement of introducing aerosol-cloud interactions is remarkable with respect to the case of including just aerosol-radiation effects. During the dust episode ( Fig. 1 right column) the analysis of the results is similar as for the fires case (averaged-domain underestimations around -1.0 K in DE-C11 to -0.56 K in ES-C21). Here the inclusion of ARI (C22) leads to a mean increase of the bias of +10.2%, but ARI+ACI (C23) leads to a very limited improvement of the simulations with respect to the base case (C21), generally reductions of the bias around -0.4%.

A similar discussion can be made for mean temperature. During the fires episode (left column of Fig. 2) all runs (but DE3) tend to underestimate the domain-averaged mean temperature (biases ranging from -0.4 K in ES1-C11 to +1.0 in DE3-C11). Here, the ensemble (ENS) simulation clearly outperforms the individual simulations (bias of -0.2 K in ENS-C11). Again, the model skill does not improve for mean temperature when including ARI or ARI+ACI (bias increase by 46.0% and 56.2%, respectively for the fires episode) but in the case of DE3-C12 simulation (including ARI reduces the bias by -27.3%). During the dust episode (right column of Fig. 2), there is a general averaged overestimation of mean



temperature (+0.4 in ES1-C21 to 0.8 K in DE3-C21). Conversely to the fires episode, the inclusion of ARI and ARI+ACI improves the bias (reductions of this variable of -13.4% in C22 and -4.2% in C23). The reduction of the bias when including ARI+ACI is especially remarkable for the ensemble of simulations, where the bias decreases by -24.4% in ENS-C23.

Last, minimum temperature during the fire episodes is shown in the left column of Fig. 3. Here results are very different to analyze for improvements or worsening of the bias, since the domain-averaged errors are in the order of -0.01 K for WRF-based models in C11 and C12, so a very slight difference would lead to a percentage increase (or reduction) of the bias compared to the base case. However, for DE3-C11 the bias is larger (up to +1.6 K for minimum temperature averaged over all the domain) and the inclusion of ARI leads only to a small improvement (-1.5%). The dust case (right column of Fig. 3) shows a general overestimation of minimum temperature, with base-case biases ranging from +0.5 K in ES1-C21 to +1.8 K in DE3-C21. Here, the inclusion of ARI and ARI+ACI slightly improves the bias (reductions of -10.5% in C22 and -5.0% in C23). Here again, the improvement of the ENS-C22 and ENS-C23 simulations is larger than for the rest of the models (reductions of the bias of -29.7% and -38.2% for ARI and ARI+ACI, respectively).

### 3.2 Temporal correlation

The temporal correlation (estimated through the coefficient of determination,  $\rho^2$ ) between simulated and observed series is shown in Fig. 4, 5 and 6 for mean maximum and mean minimum temperature, in that order. The first column in each panel represents the value of  $\rho^2$  of the base case (C11 or C21) of each individual model (or the ensemble) with respect to the E-OBS database. The center (C12 or C22) and right (C13 and C23) columns indicate the increase (red values) or decrease (blue value) of the  $\rho^2$  for each simulation with respect to the case not including feedbacks. Then, that gives an idea in the improvement (or not) in the skill of the model for representing the time evolution of our series when compared to the observations.



For maximum, mean and minimum temperature during the fires episode (left side of Fig. 4, 5 and 6, respectively), domain-averaged  $\rho^2$  is higher than 0.5 for all models (0.52 in CS1-C11 minimum temperature to 0.78 in DE3-C11 mean temperature). In general, coefficients of determination are highest for mean temperature (0.60 to 0.78) and lowest for minimum temperature (0.50 to 0.56), presenting the ensemble always maximum time  $\rho^2$  (0.75, 0.79 and 0.61, respectively for maximum, mean and minimum temperature). The highest  $\rho^2$  values are found over the north and west part of the domain (above 0.8 in mean temperature) and the lowest mainly over south and southeast area of the domain (under 0.2). According to the improvement with respect to C11 case, when analyzing the inclusion of the ARI and ARI+ACI a general improvement is observed for maximum and mean temperature, with positive values reaching up to 0.18 (domain-averaged values improve for individual models around 1% for maximum, 0.3% for mean temperature). Correlation with minima experiences a slight decrease (-0.4%) when including ARI or ARI+ACI for the ensemble mean.

During dust episode (right side of Fig. 4, 5 and 6), domain-averaged  $\rho^2$  is higher than for the fires episode for all models and variables in the base case (0.76 in DE3-C21 minima to 0.90 in DE3-C21 mean temperature), with the ensemble again providing the highest correlation (values ranging from 0.88 for maximum, 0.91 for mean and 0.84 for minimum temperature). As well as before, the inclusion of the ARI and ARI+ACI shows an improvement over some areas in the order of 0.17 for mean and maximum temperature, with domain-averaged improvements of 0.3% in C22-C23 for maximum temperature, and 0.2% in C22-C23 for mean temperature and 0.1% in C23 for minimum temperature, with no improvement for C22 in this latter variable).

### 3.3 Temporal variability

The results for the daily variability of maximum, mean and minimum temperature have been obtained by calculating the standard deviation of the daily mean series at each grid point of all the land grid points of the corresponding domain for the fires and dust episodes.

Considering maximum temperature, in the fires episode (left column of Fig. 7), all runs tend to slightly overestimate the standard deviation of maximum temperature for the base



case (no radiative feedbacks), with biases of maximum temperature standard deviation varying between +1.28 K for DE3-C11 to +0.25 K for ES1-C11. The biases of the standard deviation are reduced by -22.6% (on average) when including the ARI, with reductions in the biases of the standard deviation ranging from -34.2% in ES1-C12 and -8.6% for DE3-C12. For the ARI+ACI simulations the average reduction of the bias is -41.21% (-56.9% for ES1-C13 and -24.40% for CS2-C13). The rest of the models and cases show an intermediate behavior for representing the variability, with the best skills always for the cases including the ARI+ACI interactions. Analogous results can be found for maximum temperature during the dust episode (right column of Fig. 7): the inclusion of aerosol feedbacks generally improve the representation of the temporal variability of maximum temperature, with an average reduction of the bias of the standard deviation of -5.9% (-16.6%) for ARI (ARI+ACI) simulations.

For mean temperature during the fires episode, (left column of Fig. 8) all runs tend to overestimate the standard deviation for the base case (no radiative feedbacks), with biases of mean temperature standard deviation between +0.2 to +1.1 K. As for the maximum temperature, the biases of the standard deviation are reduced on -41.8% (on average) when including the ARI and -66.5% for the ARI+ACI simulations, with reductions in the biases of the standard deviation ranging from -8.5% in the DE3-C12 simulation to -78.2% in the ES1-C13 case. Similar to the maximum temperature, the rest of the models and cases show an intermediate representation the variability of the mean temperature, with the best skills always for the cases including the ARI+ACI interactions. Results for the dust episode are shown in the right column of Fig. 8. The standard deviation tends to be overestimated by all models in the north of Africa and central Europe, and underestimated in the eastern part of the target domain. Overall, the inclusion of ARI does not lead to better skills of the models when representing the temporal variability (+2.4%), and for ARI+ACI the skill improved only marginally (reductions of -0.6%).

With respect to the minimum temperature, for the fires episode (left column of Fig. 9) all runs tend to overestimate the standard deviation. Biases of the minimum temperature



standard deviation range between +0.4 K for the WRF-Chem-based simulations and +1.0 K for DE3-C11. The high-resolution CS2-C11 simulation presents the lowest bias (+0.3 K).

If considering the biases of the standard deviation, there is a slight improvement when including ARI or ARI+ACI for the fires episode, while a slight worsening is depicted for the dust case. The variations in the biases of the standard deviation are on average -2.1% and -4.9% respectively for the ARI and ARI+ACI simulations (+3.4% and +5.4% for the dust episode).

### 3.4 Spatial variability

Taylor diagrams (Taylor, 2001) allow an easy comparison between the spatial and temporal patterns of two fields (Rauscher et al., 2010). In Fig. 10 shows the relative spatial standard deviation (radial distance from the origin) and the correlation (the cosine of the angular coordinate) with E-Obs. Model results with good performance in terms of spatial variability and correlation are located closer to the standard deviation ratio 1 and correlation 1, which corresponds to E-OBS (indicated by the small black asterisk). For maximum, mean and minimum temperature, the diverse models (and configurations) show a narrow spread in the representation of the spatial structure of the standard deviation.

With respect to the mean field of maximum temperature (left column in Fig. 10) all models perform well for the fires period (top row), with high spatial correlations (over 0.9) and a normalized standard deviation close to observations. However, the no radiative feedback configuration (C11 cases in Fig. 10) represent excessive spatial variability (standard deviation ratio over 1). The spatial variability of the daily standard deviation for the ARI simulations (asterisks in Fig. 10, C12 cases), as well as for ARI+ACI simulations (squares, C13 cases) is substantially improved, despite the spatial correlation remains practically constant for all models. Since there is a positive bias in the models when representing the spatial variability in the no radiative feedbacks simulations, the inclusion of radiative effects reduces the variability and therefore improves its spatial patterns. Analogous results can be found for the dust episode (bottom row, Fig. 10), with a larger agreement between models, and lower dif-



ferences between C21, C22 and C23 cases (no feedbacks, ARI and ARI+ACI simulations, in that order).

With respect to the mean temperature (center column in Fig. 10), the models perform very similarly with each other, showing a high spatial correlation with the observations (over 0.9 for all models and cases), with a small overestimation of the spatial variability for the C11 (fire episode, no radiative feedbacks) case (top row), which improves when including the ARI and ARI+ACI interactions. Similarly, the spatial variability is slightly overestimated for the C21 (dust, no radiative feedbacks) case, except for the DE3 model. Generally, the models better capture the spatial structure of the variability during the fires and dust cases (Fig. 10, center column) when including the radiative feedbacks. The correlation is only slightly improved for the ARI and ARI+ACI cases (except for ENS simulations, which will be discussed below), and is always higher for the mean temperature than for maximum temperature.

The minimum temperature (Fig. 10, right column) is captured with quality as the maximum and mean temperature. While for the fire episode the models (in all cases) tend to provide a higher spatial variability than the observations, the spatial variability is underestimated for the dust episode, but with a high correlation (over 0.9) for both episodes. For this variable, the improvement of including the radiative feedbacks is not so evident, since the spatial variability does not generally improve for C12, C13, C22 or C23 cases with respect to the configuration without radiative feedbacks. Moreover, the correlation coefficient is even slightly reduced with the inclusion of ARI or ARI+ACI.

Last, the added value of considering the ensemble mean of all available simulations in each episode and case is clear for the fires episode, but not that obvious for the dust period. For the fire episode, the ensemble mean outperforms individual models in terms of the standard deviation and the correlation coefficient, especially for mean temperature, where the correlation increases up to 0.99 for the ENS-C11 case.

Generally the skill of most models improves when aerosol-meteorology interactions are taken into account.



For the dust case, the ensemble mean outperforms the individual models for representing the standard deviation (that is, the spatial variability). However, the spatial correlation coefficient is somewhat reduced as compared to the individual models.

#### 4 Summary and conclusions

5 This study shows a collective operational evaluation of the temperature at 2 meters (maximum, mean and minimum) simulated by the coupled chemistry and meteorology models under the umbrella of COST Action ES1004 for a wildfires and a dust episode in the year 2010. The meteorological parameters considered in this assessment are important to understand the effect of the aerosol interactions with clouds and radiation. In this sense, this  
10 study complements other several analysis (e.g. Brunner et al. (2015), Makar et al. (2015b), forkeletal2015) by analyzing whether the inclusion of the radiative feedbacks improves or not the representation of the temperature field (maximum, mean and minimum) in an ensemble of simulations.

Focusing on the bias, in both episodes there is a general underestimation of the studied variables, being most noticeable in maximum temperature. In general, there is not a  
15 straightforward conclusion with respect to the improvement (or not) of the bias when introducing aerosol radiative feedbacks. Broadly, the biases are improved when including ARI or ARI+ACI in the dust case, but no evident improvements are found for the heatwave/wildfires episode. Although the ensemble does not outperform the individual models (in general), the  
20 improvements found when including ARI and ARI+ACI are by far more remarkable for the ensemble than for the individual models.

With respect to the temporal correlation, maximum and mean temperatures in the fires and dust episode show higher correlations over most of the domain when considering C11 case with respect to the E-OBS database than minimum temperature. During these  
25 episodes, a twofold conclusion can be obtained: (1) the ensemble of simulations always outperforms the representation of the temporal variability of the series; and (2) an improve-



ment of the  $\rho^2$  coefficient is found when considering ARI or ARI+ACI feedbacks (in both episodes).

Regarding the temporal variability, during the fire episode there is a general pronounced overestimation of the standard deviation of the studied variables. Here, the inclusion of aerosol feedbacks largely improves the representation of the temporal variability of the three studied variables (reduction of the bias of the standard deviation) showing the best skills for the cases including the ARI+ACI interactions, with a reduction of bias of the standard deviation by as much as 75%. Very similar results can be found for the dust episode. Generally, it is for the temporal variability where the inclusion of the aerosol radiative feedbacks shows the largest improvements and results in an added value of the computational effort made to include direct aerosol radiation interactions and aerosol cloud interactions in the models. Last, with respect to the spatial variability for maximum and mean temperature, the inclusion of radiative effects reduces the variability and improves the spatial patterns for both episodes. For the minimum temperature, the improvement of including the radiative feedbacks is less evident.

In order to further investigate the impact of including the aerosol interactions in online coupled models, episodes with stronger effects on the aerosol cloud interactions should be considered since the selected episodes during EuMetChem Cost Action were mainly related to ARI. Moreover, during the dust episode, most of the ARI+ACI differences found in the models with respect to the base case were found over the Mediterranean sea, but the observational data E-OBS only has values over land. Unfortunately part of the interpretation of the results may be missed due to the unavailability of this database over the ocean. Furthermore, it must be considered, that all results for the ARI+ACI cases were from WRF-Chem simulations.

There are still modelling issues regarding the representation of the field of temperature, where maximum temperatures are underestimated and minimum temperatures are overestimated and the inclusion of the aerosol feedbacks does not improve this situation. Nevertheless, in this study, a general improvement of the temporal variability and correlation has been seen. These improvements may be important not only for certain episodes, as



analyzed here, by also for the representation of the climatology of temperatures. However, climatic-representative periods should be covered in further studies.

*Acknowledgements.* We acknowledge the E-OBS dataset from the EU-FP6 project ENSEMBLES (<http://ensembles-eu.metoffice.com>) and the data providers in the ECAD project (<http://www.ecad.eu>).

5 This research have been carried out under the support of the Project REPAIR (CGL2014-59677-R), funded by the Spanish Ministerio de Economía y Competitividad (MINECO) and the FEDER European program. Rocío Baró acknowledges the FPU scholarship (Ref. FPU12/05642) by the Spanish Ministerio de Educación, Cultura y Deporte. Pedro Jiménez-Guerrero acknowledges the fellowship 19677/EE/14 of the Programme Jiménez de la Espada de Movilidad, Cooperación e Internacionalización (Fundación Séneca-Agencia de Ciencia y Tecnología de la Región de Murcia, PCTIRM 10 2011-2014). Finally, we acknowledge the support of the European groups through COST Action ES1004 EuMetChem and the Air Quality Modelling Evaluation International Initiative (AQMEII) and the EuMetChem COST ACTION ES1004 have supported these works.

## References

- 15 Ackermann, I. J., Hass, H., Memmesheimer, M., Ebel, A., Binkowski, F. S., and Shankar, U.: Modal aerosol dynamics model for Europe: Development and first applications, *Atmospheric environment*, 32, 2981–2999, 1998.
- Alapaty, K., Mathur, R., Pleim, J., Hogrefe, C., Rao, S. T., Ramaswamy, V., Galmarini, S., Schaap, M., Makar, P., Vautard, R., Makar, P., Baklanov, A., Kallos, G., Vogel, B., Hogrefe, C., and Sokhi, R.: 20 New Directions: Understanding interactions of air quality and climate change at regional scales, *Atmospheric Environment*, 49, 419–421, 2012.
- Baklanov, A., Schlünzen, K., Suppan, P., Baldasano, J., Brunner, D., Aksoyoglu, S., Carmichael, G., Douros, J., Flemming, J., Forkel, R., Galmarini, S., Gauss, M., Grell, G., Hirtl, M., Joffre, S., Jorba, O., Kaas, E., Kaasik, M., Kallos, G., Kong, X., Korsholm, U., Kurganskiy, A., Kushta, 25 J., Lohmann, U., Mahura, A., Manders-Groot, A., Maurizi, A., Moussiopoulos, N., Rao, S. T., Savage, N., Seigneur, C., Sokhi, R. S., Solazzo, E., Solomos, S., Sørensen, B., Tsegas, G., Vignati, E., Vogel, B., and Zhang, Y.: Online coupled regional meteorology chemistry models in Europe: current status and prospects, *Atmospheric Chemistry and Physics*, 14, 317–398, 2014.



- Bangert, M., Kottmeier, C., Vogel, B., and Vogel, H.: Regional scale effects of the aerosol cloud interaction simulated with an online coupled comprehensive chemistry model, *Atmospheric Chemistry and Physics*, 11, 4411–4423, 2011.
- Baró, R., Jiménez-Guerrero, P., Balzarini, A., Curci, G., Forkel, R., Grell, G., Hirtl, M., Honzak, L., Langer, M., Pérez, J. L., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., and Zabkar, R.: Sensitivity analysis of the microphysics scheme in WRF-Chem contributions to AQMEII phase 2, *Atmospheric Environment*, 115, 620 – 629, 2015.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rash, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.-Y.: Clouds and Aerosols. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, United Kingdom and New York, USA, 2013.
- Brunner, D., Savage, N., Jorba, O., Eder, B., Giordano, L., Badia, A., Balzarini, A., Baró, R., Bianconi, R., Chemel, C., Curci, G., Forkel, R., Jiménez-Guerrero, P., Hirtl, M., Hodzic, A., Honzak, L., Im, U., Knote, C., Makar, P., Manders-Groot, A., van Meijgaard, E., Neal, L., Pérez, J. L., Pirovano, G., José, R. S., Schroder, W., Sokhi, R. S., Syrakov, D., Torian, A., Tuccella, P., Werhahn, J., Wolke, R., Yahya, K., Zabkar, R., Zhang, Y., Hogrefe, C., and Galmarini, S.: Comparative analysis of meteorological performance of coupled chemistry-meteorology models in the context of AQMEII phase 2, *Atmospheric Environment*, 115, 470 – 498, 2015.
- Chung, C. E.: *Aerosol direct radiative forcing: a review*, INTECH Open Access Publisher, 2012.
- Doms, G., Förstner, J., Heise, E., Herzog, H., Mironov, D., Raschendorfer, M., Reinhardt, T., Ritter, B., Schrodin, R., Schulz, J., and Vogel, G.: A description of the nonhydrostatic regional COSMO model, Part II: Physical Parameterization, <http://www.cosmo-model.org/>, 2011.
- Forkel, R., Werhahn, J., Hansen, A. B., McKeen, S., Peckham, S., Grell, G., and Suppan, P.: Effect of aerosol-radiation feedback on regional air quality. A case study with WRF-Chem, *Atmospheric Environment*, 53, 202–211, 2012.
- Forkel, R., Balzarini, A., Baró, R., Bianconi, R., Curci, G., Jiménez-Guerrero, P., Hirtl, M., Honzak, L., Lorenz, C., Im, U., Pérez, J. L., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., and Zabkar, R.: Analysis of the WRF-Chem contributions to AQMEII phase2 with respect to aerosol radiative feedbacks on meteorology and pollutant distributions, *Atmospheric Environment*, 115, 630 – 645, 2015.
- Forkel, R., Brunner, D., Baklanov, A., Balzarini, A., Hirtl, M., Honzak, L., Jiménez-Guerrero, P., Jorba, O., Pérez, J., San José, R., Schröder, W., Tsegas, G., Werhahn, J., Wolke, R., and Zabkar, R.: A



- Multi-model Case Study on Aerosol Feedbacks in Online Coupled Chemistry-Meteorology Models Within the COST Action ES1004 EuMetChem, in: Air Pollution Modeling and its Application XXIV, pp. 23–28, Springer, 2016.
- Galmarini, S., Hogrefe, C., Brunner, D., Makar, P., and Baklanov, A.: Preface, Atmospheric Environment, pp. 340–344, 2015.
- 5 Ghan, S. J. and Schwartz, S. E.: Aerosol properties and processes: A path from field and laboratory measurements to global climate models, Bulletin of the American Meteorological Society, 88, 1059–1083, 2007.
- Grell, G. and Baklanov, A.: Integrated modeling for forecasting weather and air quality: A call for fully coupled approaches, Atmospheric Environment, 45, 6845–6851, 2011.
- 10 Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled online chemistry within the WRF model, Atmospheric Environment, 39, 6957–6975, 2005.
- Guenther, A., Zimmermanharley, P., Monson, R., and Fall, R.: Isoprene and monoterpene rate variability: model evaluations and sensitive analyses, Journal of Geophysical Research, 98, 10 799–10 808, 1993.
- 15 Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P., Geron, C., et al.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), Atmospheric Chemistry and Physics, 6, 3181–3210, 2006.
- 20 Hansen, J., Sato, M., and Ruedy, R.: Radiative forcing and climate response, Journal of Geophysical Research: Atmospheres, 102, 6831–6864, 1997.
- Haylock, M., Hofstra, N., Klein Tank, A., Klok, E., Jones, P., and New, M.: A European daily high-resolution gridded data set of surface temperature and precipitation for 1950–2006, Journal of Geophysical Research: Atmospheres, 113, D20, doi:10.1029/2008JD010201, 2008.
- 25 Hong, S.-Y., Noh, Y., and Dudhia, J.: A new vertical diffusion package with an explicit treatment of entrainment processes, Monthly Weather Review, 134, 2318–2341, 2006.
- Im, U., Bianconi, R., Solazzo, E., Kioutsioukis, I., Badia, A., Balzarini, A., Baró, R., Bellasio, R., Brunner, D., Chemel, C., Curci, G., Flemming, J., Forkel, R., Giordano, L., Jiménez-Guerrero, P., Hirtl, M., Hodzic, A., Honzak, L., Jorba, O., Knote, C., Kuenen, J. J., Makar, P. A., Manders-Groot, A., Neal, L., Pérez, J. L., Pirovano, G., Pouliot, G., Jose, R. S., Savage, N., Schroder, W., Sokhi, R. S., Syrakov, D., Torian, A., Tuccella, P., Werhahn, J., Wolke, R., Yahya, K., Zabkar, R., Zhang, Y., Zhang, J., Hogrefe, C., and Galmarini, S.: Evaluation of operational on-line-coupled regional air



- quality models over Europe and North America in the context of AQMEII phase 2. Part I: Ozone, Atmospheric Environment, 115, 404–420, 2015a.
- 5 Im, U., Bianconi, R., Solazzo, E., Kioutsioukis, I., Badia, A., Balzarini, A., Baró, R., Bellasio, R., Brunner, D., Chemel, C., Curci, G., van der Gon, H. D., Flemming, J., Forkel, R., Giordano, L., Jiménez-Guerrero, P., Hirtl, M., Hodzic, A., Honzak, L., Jorba, O., Knote, C., Makar, P. A., Manders-Groot, A., Neal, L., Pérez, J. L., Pirovano, G., Pouliot, G., Jose, R. S., Savage, N., Schroder, W., Sokhi, R. S., Syrakov, D., Torian, A., Tuccella, P., Wang, K., Werhahn, J., Wolke, R., Zabkar, R., Zhang, Y., Zhang, J., Hogrefe, C., and Galmarini, S.: Evaluation of operational online-coupled regional air quality models over Europe and North America in the context of AQMEII phase 2. Part II: Particulate matter, Atmospheric Environment, 115, 421–441, 2015b.
- 10 Jiménez-Guerrero, P., Montávez, J., Domínguez, M., Romera, R., Fita, L., Fernández, J., Cabos, W., Liguori, G., and Gaertner, M.: Mean fields and interannual variability in RCM simulations over Spain: the ESCENA project, Climate Research, 57, 201–220, 2013.
- 15 Kioutsioukis, I., Im, U., Solazzo, E., Bianconi, R., Badia, A., Balzarini, A., Baró, R., Bellasio, R., Brunner, D., Chemel, C., Curci, G., van der Gon, H. D., Flemming, J., Forkel, R., Giordano, L., Jiménez-Guerrero, P., Hirtl, M., Jorba, O., Manders-Groot, A., Neal, L., Pérez, J. L., Pirovano, G., San Jose, R., Savage, N., Schroder, W., Sokhi, R. S., Syrakov, D., Tuccella, P., Werhahn, J., Wolke, R., Hogrefe, C., and Galmarini, S.: Insights into the deterministic skill of air quality ensembles from the analysis of AQMEII data, Atmospheric Chemistry and Physics, 16, 15 629–15 652, 2016.
- 20 Knutti, R., Furrer, R., Tebaldi, C., Cermak, J., and Meehl, G. A.: Challenges in combining projections from multiple climate models, Journal of Climate, 23, 2739–2758, 2010.
- 25 Kong, X., Forkel, R., Sokhi, R. S., Suppan, P., Baklanov, A., Gauss, M., Brunner, D., Baró, R., Balzarini, A., Chemel, C., Curci, G., Jiménez-Guerrero, P., Hirtl, M., Honzak, L., Im, U., Pérez, J. L., Pirovano, G., Jose, R. S., Schlünzen, K. H., Tsegas, G., Tuccella, P., Werhahn, J., Zabkar, R., and Galmarini, S.: Analysis of meteorology–chemistry interactions during air pollution episodes using online coupled models within AQMEII phase-2, Atmospheric Environment, 115, 527–540, 2015.
- 30 Kuenen, J., Visschedijk, A., Jozwicka, M., and Denier Van der Gon, H.: TNO-MACC\_II emission inventory; a multi-year (2003–2009) consistent high-resolution European emission inventory for air quality modelling, Atmospheric Chemistry and Physics, 14, 10 963–10 976, 2014.
- Landgren, O. A., Haugen, J. E., and Førland, E. J.: Evaluation of regional climate model temperature and precipitation outputs over Scandinavia, Climate Research, 60, 249–264, 2014.




- Lin, Y.-L., Farley, R. D., and Orville, H. D.: Bulk parameterization of the snow field in a cloud model, *Journal of Climate and Applied Meteorology*, 22, 1065–1092, 1983.
- Lohmann, U. and Feichter, J.: Global indirect aerosol effects: a review, *Atmospheric Chemistry and Physics*, 5, 715–737, 2005.
- 5 Makar, P., Gong, W., Milbrandt, J., Hogrefe, C., Zhang, Y., Curci, G., Zabkar, R., Im, U., Balzarini, A., Baró, R., Bianconi, R., Cheung, P., Forkel, R., Gravel, S., Hirtl, M., Honzak, L., Hou, A., Jiménez-Guerrero, P., Langer, M., Moran, M., Pabla, B., Pérez, J., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., Zhang, J., and Galmarini, S.: Feedbacks between air pollution and weather, Part 1: Effects on weather, *Atmospheric Environment*, 115, 442–469, 2015a.
- 10 Makar, P., Gong, W., Hogrefe, C., Zhang, Y., Curci, G., Zabkar, R., Milbrandt, J., Im, U., Balzarini, A., Baró, R., Bianconi, R., Cheung, P., Forkel, R., Gravel, S., Hirtl, M., Honzak, L., Hou, A., Jiménez-Guerrero, P., Langer, M., Moran, M., Pabla, B., Pérez, J., Pirovano, G., José, R. S., Tuccella, P., Werhahn, J., Zhang, J., and Galmarini, S.: Feedbacks between air pollution and weather, part 2: effects on chemistry, *Atmospheric Environment*, 115, 499–526, 2015b.
- 15 Meier, J., Tegen, I., Heinold, B., and Wolke, R.: Direct and semi-direct radiative effects of absorbing aerosols in Europe: Results from a regional model, *Geophysical Research Letters*, 39, L09802, doi:10.1029/2012GL050994, 2012.
- Morrison, H., Thompson, G., and Tatarskii, V.: Impact of cloud microphysics on the development of trailing stratiform precipitation in a simulated squall line: Comparison of one-and two-moment schemes, *Monthly Weather Review*, 137, 991–1007, 2009.
- 20 Myhre, G., Samset, B. H., Schulz, M., Balkanski, Y., Bauer, S., Berntsen, T. K., Bian, H., Bellouin, N., Chin, M., Diehl, T., Easter, R. C., Feichter, J., Ghan, S. J., Hauglustaine, D., Iversen, T., Kinne, S., Kirkevåg, A., Lamarque, J.-F., Lin, G., Liu, X., Lund, M. T., Luo, G., Ma, X., van Noije, T., Penner, J. E., Rasch, P. J., Ruiz, A., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., Wang, P., Wang, Z., Xu, L., Yu, H., Yu, F., Yoon, J.-H., Zhang, K., Zhang, H., and Zhou, C.: Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations, *Atmospheric Chemistry and Physics*, 13, 1853–1877, 2013.
- 25 Péré, J., Bessagnet, B., Mallet, M., Waquet, F., Chiapello, I., Minvielle, F., Pont, V., and Menut, L.: Direct radiative effect of the Russian wildfires and its impact on air temperature and atmospheric dynamics during August 2010, *Atmospheric Chemistry and Physics*, 14, 1999–2013, 2014.
- 30 Pouliot, G., van der Gon, H. A. D., Kuenen, J., Zhang, J., Moran, M. D., and Makar, P. A.: Analysis of the emission inventories and model-ready emission datasets of Europe and North America for phase 2 of the AQMEII project, *Atmospheric Environment*, 115, 345–360, 2015.




- Rauscher, S. A., Coppola, E., Piani, C., and Giorgi, F.: Resolution effects on regional climate model simulations of seasonal precipitation over Europe, *Climate Dynamics*, 35, 685–711, 2010.
- San José, R., Pérez, J., Balzarini, A., Baró, R., Curci, G., Forkel, R., Galmarini, S., Grell, G., Hirtl, M., Honzak, L., Im, U., Jiménez-Guerrero, P., Langer, M., Pirovano, G., Tuccella, P., Werhahn, J., Zabkar, R.: Sensitivity of feedback effects in CBMZ/MOSAIC chemical mechanism, *Atmospheric Environment*, 115, 646–656, 2015.
- Simpson, D., Fagerli, H., Jonson, J., Tsyro, S., Wind, P., and Tuovinen, J.: The EMEP Unified Eulerian Model. Model Description. EMEP MSC-W Report 1/2003, The Norwegian Meteorological Institute, Oslo, Norway, 2003.
- Soares, J., Sofiev, M., and Hakkarainen, J.: Uncertainties of wild-land fires emission in AQMEII phase 2 case study, *Atmospheric Environment*, 115, 361–370, 2015.
- Sofiev, M., Vankevich, R., Lotjonen, M., Prank, M., Petukhov, V., Ermakova, T., Koskinen, J., and Kukkonen, J.: An operational system for the assimilation of the satellite information on wild-land fires for the needs of air quality modelling and forecasting, *Atmospheric Chemistry and Physics*, 9, 6833–6847, 2009.
- Solazzo, E. and Galmarini, S.: A science-based use of ensembles of opportunities for assessment and scenario studies, *Atmospheric Chemistry and Physics*, 15, 2535–2544, 2015.
- Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *Journal of Geophysical Research: Atmospheres*, 106, 7183–7192, 2001.
- Twomey, S.: The influence of pollution on the shortwave albedo of clouds, *Journal of the Atmospheric Sciences*, 34, 1149–1152, 1977.
- Vautard, R., Moran, M. D., Solazzo, E., Gilliam, R. C., Matthias, V., Bianconi, R., Chemel, C., Ferreira, J., Geyer, B., Hansen, A. B., Jericevic, A., Prank, M., Segers, A., Silver, J. D., Werhahn, J., Wolke, R., Rao, S., and Galmarini, S.: Evaluation of the meteorological forcing used for the Air Quality Model Evaluation International Initiative (AQMEII) air quality simulations, *Atmospheric Environment*, 53, 15–37, 2012.
- Wolke, R., Schröder, W., Schrödner, R., and Renner, E.: Influence of grid resolution and meteorological forcing on simulated European air quality: a sensitivity study with the modeling system COSMO–MUSCAT, *Atmospheric environment*, 53, 110–130, 2012.
- Yang, Q., Gustafson Jr, W., Fast, J. D., Wang, H., Easter, R. C., Morrison, H., Lee, Y.-N., Chapman, E. G., Spak, S., and Mena-Carrasco, M.: Assessing regional scale predictions of aerosols, marine stratocumulus, and their interactions during VOCALS-REx using WRF-Chem, *Atmospheric Chemistry and Physics*, 11, 11 951–11 975, 2011.




Zaveri, R. A., Easter, R. C., Fast, J. D., and Peters, L. K.: Model for simulating aerosol interactions and chemistry (MOSAIC), *Journal of Geophysical Research*, 113, D13204, doi:10.1029/2007JD008782, 2008.




**Figure 1.** (Top row) Maximum temperature (TMAX) for the fires (left) and dust (right) episodes, as derived from E-OBS database (in K). The panel below represents the bias for the fires (left) and dust (right) episodes of each simulation with respect to the E-OBS database. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.




**Figure 2.** (Top row) Mean temperature (TEMP) for the fires (left) and dust (right) episodes, as derived from E-OBS database (in K). The panel below represents the bias for the fires (left) and dust (right) episodes of each simulation with respect to the E-OBS database. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.




**Figure 3.** (Top row) Minimum temperature (TMIN) for the fires (left) and dust (right) episodes, as derived from E-OBS database (in K). The panel below represents the bias for the fires (left) and dust (right) episodes of each simulation with respect to the E-OBS database. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.




**Figure 4.** (Top row) Time determination coefficient ( $\rho^2$ ) (model vs. E-OBS) of the maximum temperature (TMAX) for the fires (left panel) and dust (right panel) episodes. The first column in each panel below represents the value of  $\rho^2$  of the no radiative feedback case with respect to the E-OBS database. The center and right columns indicate the increase (red values) or decrease (blue value) of each simulation with respect to the case not including feedbacks. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.




**Figure 5.** (Top row) Time determination coefficient ( $\rho^2$ ) (model vs. E-OBS) of the mean temperature (TEMP) for the fires (left panel) and dust (right panel) episodes. The first column in each panel below represents the value of  $\rho^2$  of the no radiative feedback case with respect to the E-OBS database. The center and right columns indicate the increase (red values) or decrease (blue value) of each simulation with respect to the case not including feedbacks. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.




**Figure 6.** (Top row) Time determination coefficient ( $\rho^2$ ) (model vs. E-OBS) of the minimum temperature (TMIN) for the fires (left panel) and dust (right panel) episodes. The first column in each panel below represents the value of  $\rho^2$  of the no radiative feedback case with respect to the E-OBS database. The center and right columns indicate the increase (red values) or decrease (blue value) of each simulation with respect to the case not including feedbacks. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.




**Figure 7.** (Top row) Standard deviation (STD) of the maximum temperature (TMAX) for the fires (left) and dust (right) episodes, as derived from E-OBS database (in K). The panel below represents the bias for the standard deviation of the fires (left) and dust (right) episodes of each simulation with respect to the E-OBS database. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.



**Figure 8.** (Top row) Standard deviation (STD) of the mean temperature (TEMP) for the fires (left) and dust (right) episodes, as derived from E-OBS database (in K). The panel below represents the bias for the standard deviation of the fires (left) and dust (right) episodes of each simulation with respect to the E-OBS database. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.



**Figure 9.** (Top row) Standard deviation (STD) of the minimum temperature (TMIN) for the fires (left) and dust (right) episodes, as derived from E-OBS database (in K). The panel below represents the bias for the standard deviation of the fires (left) and dust (right) episodes of each simulation with respect to the E-OBS database. NRF: no radiative feedbacks; ARI: aerosol-radiation interactions; ARI+ACI: as ARI including aerosol-cloud interactions.





**Figure 10.** Taylor diagrams for (left) maximum temperature, (center) mean temperature, and (right) minimum temperature for the simulations included in the analysis. The top row represents the Taylor diagrams for the fires episode, while the bottom row stands for the dust episode. The cases included are: no radiative feedbacks (filled circle), ARI (asterisk) and ARI+ACI (empty squares). Each configuration is shown in a different color: CS1 (green), CS2 (dark blue), DE3 (red), ES1 (yellow), ES3 (pink) and ENS (black).



**Table 1.** Modelling systems participating and their contributions to the case studies

	CS1	CS2	DE3	ES1	ES3
Lead Institution	UL, KIT/IMK-IFU*	UL, KIT/IMK-IFU*	IFT Leipzig	U. Murcia	UPM-ESMG
Model	WRF-Chem	WRF-Chem	COSMO-MUSCAT	WRF-Chem	WRF-Chem
Episode	Fire, Dust	Fire	Fire, Dust	Fire, Dust	Fire, Dust
Runs	NRF, ARI, ARI+ACI	NRF, ARI, ARI+ACI	NRF, ARI	NRF, ARI, ARI+ACI	NRF, ARI, ARI+ACI
Resolution	23 km	9.9 km	0.125 deg.	23 km	23 km
Microphysics	Morrison	Morrison	Kessler-type bulk	Lin	Morrison
SW Radiation	Goddard	Goddard	$\delta$ -2-stream	Goddard	RRTMG
LW Radiation	RRTM	RRTM	$\delta$ -2-stream	RRTM	RRTMG
PBL/turbulence	YSU	YSU	Prognostic TKE	YSU	YSU
Biogenic model	MEGAN (Guenther et al., 2006)	MEGAN	Guenther et al. (1993)	MEGAN	MEGAN
Gas phase	RADM2 mod.	RADM2 mod.	RACM-MIM2	RADM2	CBMZ
Aerosol	MADE/SORGAM	MADE/SORGAM	Simpson et al. (2003)	MADE/SORGAM	MOSAIC 4 bins
Model reference	Grell et al. (2005); Forkel et al. (2015)	Grell et al. (2005); Forkel et al. (2015)	Wolke et al. (2012)	Grell et al. (2005)	Grell et al. (2005)

\*Joint effort, also including ZAMG, RSE, UPM-ESMG