

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Separating agricultural and non-agricultural fire seasonality at regional scales

B. I. Magi¹, S. Rabin², E. Shevliakova², and S. Pacala²

¹University of North Carolina at Charlotte, Geography and Earth Sciences Department, Charlotte, North Carolina, USA

²Princeton University, Department of Ecology and Evolutionary Biology, Princeton, New Jersey, USA

Received: 12 April 2012 – Accepted: 22 April 2012 – Published: 11 May 2012

Correspondence to: B. I. Magi (brian.magi@uncc.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The timing and length of burning seasons in different parts of the world depend on climate, land cover characteristics, and human activities. In this study, global fire data from satellite-based instruments are used in conjunction with global gridded distributions of agricultural land cover types (defined as the sum of cropland and pasture area) to separate the seasonality of agricultural burning practices from that of non-agricultural fire. The results presented in this study show that agricultural and non-agricultural land experience broadly different fire seasonality patterns that are not always linked to climate conditions. We highlight these differences on a regional basis, examining variations in both agricultural land cover and associated cultural practices to help explain our results. While we discuss two land cover categories, the methods can be generalized to derive seasonality for any number of land uses or cover types. This will be useful as global fire models evolve to be fully interactive with land use and land cover change in the next generation of Earth system models.

1 Introduction

Fire is a phenomenon that connects the land, the atmosphere, and the climate with human behavior in many parts of the world. Global fire models rely on climate and vegetation conditions to determine flammability and fire behavior (e.g., Arora and Boer, 2005) and now explicitly consider human population density as a source of both fire ignitions and fire suppression (e.g., Pechony and Shindell, 2009). The knowledge of spatial and temporal fire patterns has improved over the past 10–15 yr with the development of satellite-based remote sensing methods to characterize fires at a global scale (e.g., Justice et al., 2002; Giglio et al., 2006a) and much finer scales (e.g., Morton et al., 2008; McCarty et al., 2009). Such remote sensing data are critical for parameterizing and validating fire models. However, remote sensing data products do not attribute observed fires to a specific cause, while fire models operate within a framework that

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



must consider underlying land cover conditions to determine the fuel load (biomass) available to burn.

Fires can be ignited by lightning, but most are a result of human practices. These include a wide range of managed activities, such as preparing cropland for planting or harvest, managing pasture for livestock or game, clearing forest for agriculture, or even unmanaged practices such as arson, accidental ignitions, or unintentional spread of managed fires into unmanaged land (Laris and Wardell, 2006; McCarty et al., 2009). Fire suppression tactics add an additional dimension to the human component of fires. Altogether, human intervention has altered the seasonality of fires in many parts of the world, both in terms of timing and magnitude (Le Page et al., 2010; Aldersley et al., 2011). A critical component of this realization is that, as suggested by Le Page et al. (2010), human burning practices do not necessarily follow the climate-driven burning season. To more completely model fires on a global scale, particularly under projected climate change, the relationship between humans, climate, land cover, and fire needs to be better characterized.

The work presented in this study quantifies the influence of agricultural burning practices on the seasonal timing of fires. We present a method that uses satellite observations of the number of fires and burned area in conjunction with global estimates of cropland and pasture extent to attribute fires and burned area to either agricultural or non-agricultural land cover. We focus on fourteen regions around the world to investigate the fire patterns unique to each.

2 Methods

2.1 Regions

We used the fourteen regions discussed by Giglio et al. (2006b) to isolate areas of the world with similar seasonal fire patterns. Table 1 lists the names of these regions along with the shorthand notation that will be used throughout this paper, and they are shown

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in Fig. 1. These regions have been used by a number of global studies of fire patterns (e.g., Giglio et al., 2010; Kloster et al., 2010; van der Werf et al., 2010; Aldersley et al., 2011).

2.2 Fire data

5 Monthly fire counts at 0.5° resolution are taken from the Moderate Resolution Imaging Spectroradiometer (MODIS) Collection 5 Climate Modeling Grid (CMG) fire count data product (Giglio et al., 2006a). Both the Terra MODIS and Aqua MODIS fire counts from the beginning of their datasets (November 2000 and July 2002, respectively) through December 2009 are considered in this study. The processing algorithm of the MODIS
10 data is discussed by Justice et al. (2002), and comparisons between Terra MODIS and Aqua MODIS are described by Giglio et al. (2006a).

Monthly burned area data at 0.5° resolution are taken from the Global Fire Emissions Database (GFED) version 3 (Giglio et al., 2010). The years 2000–2009 are considered in this study. The GFED processing algorithm (Giglio et al., 2010) uses MODIS fire counts, surface reflectances, and land-cover characteristics (Giglio et al., 2009) to produce a burned area data set that characterizes burns as small as 1.2 km^2 . Although
15 the GFED data partially depend on MODIS fire counts, we include them in this analysis because burned area is important when quantifying fire impacts on the carbon cycle.

Henceforth, we use “fire data” to refer to fire counts or burned area as appropriate,
20 but we generally focus on burned area.

2.3 Land-use data

Cropland and pasture area are taken from the History Database of the Global Environment (HYDE) version 3.1 (Klein-Goldewijk et al., 2011), which is available at $5'$ spatial resolution. In this study, we aggregate the HYDE data to 0.5° resolution to match the
25 resolution of the fire data described above, where we define total agricultural area as the sum of cropland and pasture area. The remaining land area is considered to be

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



non-agricultural. The HYDE data set is available for 2000 and 2005; we assume that the distribution of agricultural area for every month from 2000–2004 is equal to the 2000 distribution, and that the distribution for 2005–2009 is equal to that for 2005.

2.4 Data analysis

To best understand the qualitative seasonality of agricultural and non-agricultural fires, the fire data time series (count or burned area) in each grid cell is normalized to its maximum value. Every grid cell, therefore, varies between zero and one regardless of the absolute magnitude of fire counts or burned area in a particular grid cell. This normalization overcomes the issue of a region's fire counts or burned area patterns being dominated by intense burning in one part of the region, allowing regions with even minor burning to be considered in our investigation of seasonality. In this study, we refer to the processed data sets as the “normalized fire counts” and “normalized burned area”, or collectively as the “normalized fire data”.

The relationship between fire counts and GFED burned area depends on the density and moisture of the vegetation being burned (Giglio et al., 2006a). As such, the question might be raised of whether it is worthwhile to analyze these two types of data together. The graphs of regionally-averaged normalized fire data shown in Fig. 2 suggest that, regardless of the lack of direct correlation between fire counts and burned area, overall seasonality in fires is similar among the three data sets. Pearson's product-moment correlation coefficient (r) between Terra and Aqua MODIS fire counts in 2000–2009 is 0.95, and Terra MODIS fire count and GFED burned area data are similarly correlated ($r = 0.91$; Fig. S1 in the Supplement).

Using the normalized fire data sets, we examine regional subsets (Fig. 1) of the global fire data using the method described below.

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.5 Attribution method

For each region, we use a method that is based on a least-squares analysis to determine fire activity with respect to agricultural and non-agricultural land-use. In every grid cell i , we assume that

$$p_{a,i} + p_{n,i} = 1 \quad (1)$$

where $p_{a,i}$ and $p_{n,i}$ are the fractions of land area in grid cell i that are agricultural and non-agricultural, respectively. The expected amount of fire in grid cell i (E_i) is defined as

$$E_i = p_{a,i}\bar{F}_a + (1 - p_{a,i})\bar{F}_n \quad (2)$$

where \bar{F}_a and \bar{F}_n represent the regionally-averaged (i.e., over all N grid cells in the region) expected value of fire data for agricultural and non-agricultural fractions, respectively. For one month and two land-use types, we find the sum of squared differences (S) between E_i and the observed amount of fire in each grid cell i as

$$S = \sum_i^N [E_i - D_i]^2 \quad (3)$$

where D_i represents the observed fire data (count or burned area). The unknowns in the equation are \bar{F}_a and \bar{F}_n . Our goal is to find values for these unknowns that minimize S . Essentially, we are unpacking the land-use specific signals from the fire data.

To solve for the two unknowns (\bar{F}_a and \bar{F}_n), we set the partial derivative of S with respect to \bar{F}_n to zero such that

$$\frac{\partial S}{\partial \bar{F}_n} = \sum_i^N 2[1 - p_{a,i}][E_i - D_i] = 0 \quad (4)$$

This can be simplified to:

$$\bar{F}_a = \frac{\bar{D}}{\bar{p}_a} + \left(1 - \frac{1}{\bar{p}_a}\right)\bar{F}_n \quad (5)$$

where \bar{p}_a is the mean agricultural fraction and \bar{D} is the mean observed fire counts or burned area. Taking the partial derivative with respect to \bar{F}_a gives:

$$\frac{\partial S}{\partial \bar{F}_a} = \sum_i^N 2p_{a,i}[E_i - D_i] = 0 \quad (6)$$

Using Eq. (5), this can be simplified to:

$$\bar{F}_n = \bar{D} - \frac{\sigma_{p_a,D}}{\sigma_{p_a}^2} \bar{p}_a \quad (7)$$

where $\sigma_{p_a}^2$ is the variance of p_a and $\sigma_{p_a,D}$ is the covariance of p_a and D . \bar{F}_n is calculated explicitly from Eqs. (7), and (5) can then be used to determine \bar{F}_a .

When the calculation summarized by Eqs. (5) and (7) return values of $\bar{F}_a < 0$ or $\bar{F}_n < 0$, we apply the following conditional simplification to preserve a realistically constrained value of \bar{F}_a and \bar{F}_n . If $\bar{F}_a < 0$, we set $\bar{F}_a = 0$ and re-calculate \bar{F}_n from Eq. (5). If $\bar{F}_n < 0$, we set $\bar{F}_n = 0$ and re-calculate \bar{F}_a from Eq. (5).

We apply the method every month to the normalized fire data in every study region. Land-use distributions are nearly constant from 2000–2009, so the monthly variations in \bar{F}_a and \bar{F}_n are driven by variability in the monthly values of the fire data over the periods considered

2.6 Validation of the attribution method

Under ideal conditions, we would test the above method by examining fire data that we know correspond to agricultural and non-agricultural burning. This information is, however, the unknown for which we developed the method. We instead test it by masking the normalized fire data for two cases. In one case, all grid cells with non-agricultural fraction less than 0.8 are excluded; the fire data of the remaining grid cells are assumed to represent non-agricultural fire seasonality. In the second case, all grid cells

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with agricultural fraction less than 0.8 are excluded, with the fire data for the remaining grid cells assumed to represent agricultural fire seasonality. We chose 80% fractional coverage as the threshold based on the land-cover classification scheme for cropland discussed by Hansen et al. (2000). We will refer to this as the “subset method.” Table 2 shows the fraction of grid cells in that qualify for each case, and the total number of grid cells in each region.

We examine each of the 14 regions separately, determining the mean fire seasonality for the grid cells remaining after applying the threshold. As shown in Table 2, the mean fraction of non-agricultural grid cells in a region is 0.34 ± 0.25 , while the mean fraction of agricultural grid cells is 0.13 ± 0.11 . These subsets of all grid cells in a region significantly reduce the sample size for regions such as AUST, EQAS, and NHSA, which is a limitation of the subset method.

For each fire data set, the mean fire seasonality value for each month and region is calculated as the mean of the seasonality value in that month for all N grid cells in the region over the ten years in the time series. There are a total of 168 mean fire seasonality values (12 months and 14 regions) for each of the six combinations of three fire data sets and two land cover types. This is performed for seasonality values obtained using the attribution method described by Eqs. (5) and (7) and the subset method. Our goal is to test the attribution method.

3 Results and discussion

The correlation coefficients between the attribution method and the subset method results are always greater than 0.91, which is significant with $p < 0.05$. We conclude that the attribution method is an effective means to estimate fire seasonality on different land cover types. From this point forward, we only refer to results from the attribution method (summarized by Eqs. 5 and 7).

Regional fire seasonality determined from the attribution method (Fig. 3) shows that for the majority of the study regions, the agricultural fire season differs from the

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



non-agricultural fire season. These differences are broadly consistent with past studies of the human and physical drivers of global fires (e.g., Korontzi et al., 2006; Le Page et al., 2010).

To facilitate comparison of when fire is expected to occur and when it does occur, we have overlaid mean monthly flammability and lightning flash rate onto Fig. 3. Flammability is calculated using formulations described by Pechony and Shindell (2009) and depends on temperature, relative humidity, and rainfall rate. Values of flammability close to one in Fig. 3 suggest a high susceptibility to fire ignitions provided fuel (vegetation) is available to burn and there is an ignition source (i.e., humans or lightning). Values close to zero indicate that ignition is unlikely. Temperature and relative humidity data are from the National Center for Environmental Prediction reanalysis product (Kalnay et al., 1996; downloaded from <http://www.esrl.noaa.gov/psd/>), while rainfall rate data are from the Global Precipitation Climatology Project version 2.2 (Huffman et al., 2009). Lightning flash rate is a combined data product based on observations from the NASA Lightning Imaging Sensor and Optical Transient Detector (Christian et al., 2003; downloaded from <http://thunder.nsstc.nasa.gov/data/>).

Figure 4 shows maps of (a) the agricultural fraction of the land area in a grid cell and (b) the fraction of agricultural area in a grid cell that is cropland as opposed to pasture. These data represent averaged HYDE data (Klein Goldewijk et al., 2011) from 2000 and 2005. Agricultural fractions (Fig. 4a) are high throughout the world, with the exceptions being deserts, tropical rainforests, mountainous regions, and the boreal forests. Of the agricultural fraction of land, pasture is the dominant land-use type. The exceptions are the crop belts of the world, as discussed by Leff et al. (2004).

Below, we discuss the agricultural and non-agricultural fire seasonality for each of the 14 regions (Fig. 1), with section heads representing large multi-region areas. The sections are ordered from highest to lowest average annual burned area (Fig. 2, or Table 2 in Giglio et al., 2010). Generally, our results show that burning on non-agricultural lands depends more on weather than agricultural burning. The timing of agricultural

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



burning depends on the needs of farmers and pastoralists (e.g., Vigilante et al., 2004; Laris and Wardell, 2006; McCarty et al., 2009), which we discuss on a per-region basis.

3.1 Sub-Saharan Africa (NHAF, SHAF)

The non-agricultural fire season in Northern Hemisphere Africa (NHAF) begins around November–December and lasts through February, with a peak in December–January, corresponding roughly to the first half of a period of increased flammability (Fig. 3). The agricultural fire season peaks at the same time, but begins about two months earlier in the dry season (Fig. 3). Previous work has shown that most of NHAF exhibits a fire season earlier than would be expected based on an index of fire weather alone (Laris and Wardell, 2006; Le Page et al., 2010), suggesting that earlier, less weather-dependent agricultural burning has influenced the overall fire seasonality in NHAF. Laris and Wardell (2006) discuss considerable cultural and physical evidence that early dry season fire is the dominant agricultural burning practice in the West African nations of Mali and Burkina Faso, mainly because fires are easier to control and trees are less likely to be killed if vegetation is burned before the peak of the dry season. Most agricultural land in this region is pasture (Fig. 4), and Korontzi et al. (2006) showed that cropland burning accounted for only about 2% of the MODIS-observed fires from 2001–2003, while savanna fires accounted for about 75%. Results in this study thus agree with past studies and indicate that early dry season burning of agriculture – mostly pasture – is the dominant cause of fires in NHAF.

In Southern Hemisphere Africa (SHAF), we found that the non-agricultural fire season lasts from about May to October, while the agricultural fire season lasts from June to November with a smaller peak around March to April. The non-agricultural fire season in SHAF has the same shape as the agricultural fire season but is shifted about two months earlier, closely following increasing flammability (Fig. 3). Korontzi et al. (2006) found that cropland fires in SHAF occurred from about June to October and that August was the month of peak burning, which is not consistent with the peak value of agricultural burning from our results. However, cropland fires in SHAF contribute even

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



less to burning than in NHAf, comprising only 1 % of all fires while savanna fires make up 80 % (Korontzi et al., 2006).

Like NHAf, SHAF agricultural land cover is mainly pasture (Fig. 4). Le Page et al. (2010) showed that across SHAF, the month of peak flammability ranged from July–August in the northwest (Northern Angola) to November–December in the east and southeast (Mozambique, South Africa). The agricultural burning occurs in the latter half of a period of high flammability while lightning activity is increasing (Fig. 3). Le Page et al. (2010) also showed that SHAF experienced delayed overall burning (i.e., delayed with respect to flammability) along the southeastern coast; that region has a higher fraction of cropland than the rest of SHAF (Fig. 4), and this delay may partially account for our result of delayed agricultural burning. This suggests that the agricultural fire timing across SHAF is mostly due to burning practices in cropland; however, it is unclear why late dry season agricultural burning practices would be dominant in SHAF when early dry season agricultural burning practices are the norm in NHAf.

3.2 Latin America (CEAM, NHSA, SHSA)

Agricultural and non-agricultural burning in Central America (CEAM) both peak around April–May; however, the agricultural fire season begins earlier – in November–December, coincident with an increase in flammability (Fig. 3) – and ends later. In Mexico, Korontzi et al. (2006) found peak cropland burning in April–May, which they attribute to the harvest of maize, rice, sugarcane, and winter wheat. Since Mexico makes up much of the land area in CEAM and much agriculture in this region is comprised of croplands (Fig. 4), it is not surprising that we got a similar finding for CEAM’s agricultural fire season there.

The beginning of the non-agricultural fire season closer to the period of peak flammability in CEAM suggests that non-agricultural burning depends on fuel having desiccated over much of the dry season, whereas agricultural burning takes place throughout the dry season (Fig. 3). Le Page et al. (2010) found delayed burning throughout most of CEAM, which our analysis suggests is non-agricultural (Fig. 3).

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In Northern Hemisphere South America (NHTA), agricultural and non-agricultural burning are similar from January to August, but diverge from September through November. Non-agricultural lands experience a secondary peak in fires in October, while agricultural lands see a pronounced dip (Fig. 3). The February–March peak shared by agricultural and non-agricultural lands corresponds to a period of high flammability, while the secondary agricultural burning peak in October appears to be related to a peak in lightning frequency (Fig. 3). The agricultural fire seasons in CEAM and NHTA are similar, which corresponds to similar crop types between the two regions (Leff et al., 2004).

The agricultural and non-agricultural fire seasons in Southern Hemisphere South America (SHSA) correspond closely to trends in flammability. This agrees with findings by Le Page et al. (2010), which show that most of tropical SHSA exhibits no sign of a human impact on the fire season, although there are scattered zones of delayed fire seasonality. The agricultural fire season peaks around August–September (Fig. 3). This agrees well with previous work on cropland fire activity by Korontzi et al. (2006), who found peak burning late in the dry season. However, those authors also show that cropland burning accounts for only 2 % of all fires in SHSA, while fire in grassland and savanna (where most pasture is located) accounts for about 50–60 %. This suggests that pasture might dominate the agricultural fire signal, but it happens that burning in pasture-rich regions of Brazil and Argentina tends to be similarly timed (Le Page et al., 2010). Our results agree with previous work suggesting that early dry season burning in this part of the region is limited (Le Page et al., 2010), unlike the heavily pastoral NHTA.

3.3 Southeast Asia, Australia, and New Zealand (SEAS, EQAS, AUST)

In Southeast Asia (SEAS), most agricultural burning takes place in April and May, with a more minor peak around October. Agriculture in this region is dominated by croplands, most of which are concentrated in India and Bangladesh (Fig. 4). Previous analyses of agricultural fire seasonality in India have found similar patterns. Korontzi

et al. (2006) showed that most cropland fires occurred from April to June, which those authors associated with the sorghum harvest in Central India. Le Page et al. (2010) also found April–May and October–November peaks in overall burning for an especially cropland-rich part of Northern India, which they attributed to the wheat and rice harvests, respectively. Thus, it appears that agricultural burning practices in India (or practices that are similarly timed) dominate the agricultural fire signal for the entire region. This is distinct from the non-agricultural fire season, which lasts from about January to April.

The peaks of the agricultural and non-agricultural fire seasons in equatorial Asia (EQAS) are similar, but the former begins around May while the latter begins closer to July (Fig. 3). Both peak in August, which corresponds to maximum flammability (Fig. 3). However, Le Page et al. (2010) found that the fire season throughout most of this region was delayed by at least 30 days compared to flammability conditions. Most likely, the differences between our studies are an artifact of slight differences in calculation and interpretation of flammability. For example, we include rainfall rate in our flammability calculation (per Pechony and Shindell, 2009) whereas Le Page et al. (2010) do not. Rainfall rates are relatively high in EQAS (not shown), and as such this difference in method might be important. Furthermore, while our methods seem to have resulted in similar timing of peak seasonality across the region as a whole, Le Page et al. (2011) did not consider peak flammability but rather the middle of a “season” of high flammability.

Analysis of fires in Australia and New Zealand (AUST) reveals two similarly strong agricultural fire seasons, one peaking in April and the other in November. The April peak corresponds to a period of somewhat elevated flammability, while the November peak in burning occurs as flammability nears its minimum after a September peak (Fig. 3). Korontzi et al. (2006) found that cropland burning in Southern Australia peaks from March to May, which corresponds to the agricultural fire season that we estimated. However, Australia is characterized by extensive pasture (Fig. 4), and thus agricultural burning is probably dominated by pastoral management practices – indeed, the

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



number of cropland fires here is very small compared to those in savannas or grasslands (Korontzi et al., 2006). Vigilante et al. (2004) discuss the complex fire management of Northern Australian pastures and point out that while early dry season burning practices (March to May) are certainly common, late dry season fire use (October to November) is also extensive. This latter custom can be attributed mainly to Aboriginal pastoralists who prefer to wait for the dense grasses on fertile soils to fully dry before igniting fires that, as a result, are larger and more intense (Vigilante et al., 2004). Our results in AUST agree, providing strong evidence of extensive late dry season agricultural burning that is mainly due to pasture fires.

The non-agricultural fire season in AUST shows a pronounced dip in April that unexpectedly corresponds to a small peak in flammability (Fig. 3). However, the non-agricultural peak does come during the late dry season, when lightning frequency increases and vegetation is still relatively flammable (Fig. 3). Overall, the main non-agricultural fire season has the same shape as the agricultural fire season, but is shifted one month earlier. They also differ in that the non-agricultural burning dip corresponds to one of the agricultural burning peaks (Fig. 3).

3.4 Boreal and Central Asia (BOAS, CEAS)

Agricultural burning in BOAS takes place in two different seasons, with a major peak around April to May and a minor peak from August to October (Fig. 3). The first peak corresponds to a period of elevated flammability, and the second occurs as flammability approaches its minimum (Fig. 3). This fits well with a previous finding that cropland fires occur in April–May with a second peak occurring in August–October (Korontzi et al., 2006), bracketing the period of maximum flammability in the region (Fig. 3). On the other hand, non-agricultural burning in BOAS (May to September with a July peak) corresponds very closely to the period of maximum flammability and lightning activity (June to August; Fig. 3).

Like BOAS, CEAS experiences a biennial agricultural fire season, with a major peak around August and a minor peak around April (Fig. 3). Korontzi et al. (2006) found

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



similar timing and relative magnitude for cropland fires in Ukraine (part of CEAS). Although those authors found that cropland fire seasons in the broader CEAS region that were similarly timed but opposite in magnitude, the fact that our results for agricultural burning in CEAS as a whole match those for Ukraine suggests that cropland fires in that country dominate the overall agricultural signal across the region. Indeed, Korontzi et al. (2006) attributed about 30 % of the fires in CEAS to cropland fires,

Non-agricultural fire in CEAS shows a very different pattern, peaking around March–April and July–August although such burning does take place at some level throughout the year (Fig. 3). We would expect non-agricultural fire seasonality to be closely tied to the peaks of flammability and lightning activity from June to August, but our results suggest this is not the case. Le Page et al. (2010) found the middle of the flammability and overall fire seasons to vary widely across this region, the latter from February in Southern China to October in Kazakhstan. A more detailed analysis of the spatial distribution of burning and agriculture within CEAS might help explain the timing of non-agricultural burning in different parts of the region.

3.5 North America (BONA, TENA)

Agricultural burning peaks in April and May for temperate and boreal North America, respectively (TENA and BONA; Fig. 3). This is consistent with an April to June peak in North American cropland fires identified by Korontzi et al. (2006). However, this burning takes place about 2–3 months earlier than the peak period of flammability and lightning activity (Fig. 3). Our results also show evidence of a lower-amplitude but broader peak in agricultural burning from July to October, which previous researchers have associated with crop harvest (Korontzi et al., 2006; McCarty et al., 2009). Some of this burning takes place at the end of the high flammability season, but it extends well into a period of low flammability (Fig. 3). However, the strength of the first agricultural fire peak corresponds well to previous observations that the overall fire season in much of the cropland belt that extends from the Ohio River Valley in the United States to Southern Canada occurs in April to May, about 50–150 days earlier than would be expected

based on weather conditions alone (Le Page et al., 2010). Thus, our results support the idea that agricultural burning practices depend more on crop cycles than flammability (McCarty et al., 2009).

The non-agricultural fire seasons in North America last from about June to October, with a July peak in BONA and an August peak in TENA (Fig. 3). These are quite distinct from those regions' agricultural fire seasons, and are consistent with work finding that the boreal forests of Canada are most vulnerable to fire during the warmer summer months during which flammability and lightning frequency are at their maximum (Turetsky et al., 2010; Fig. 3). However, the timing we found for the non-agricultural fire season does not exactly match that of Le Page et al. (2010), who calculated it as later than July, and even as late as January, in many parts of BONA.

3.6 Europe and Middle East (EURO, MIDE)

The agricultural and non-agricultural fire seasons in Europe (EURO) are similar in terms of timing of minimum and maximum activity, but non-agricultural burning climbs steadily from January through its peak whereas agricultural burning experiences a minor peak in April followed by a dip and then a sharp increase moving into July (Fig. 3). Mixed results in this region for the timing of overall burning relative to flammability (Le Page et al., 2010) supports our finding of similarly-timed fire season. Cropland comprises 50 % or more of agricultural area in most of Europe (Fig. 4), so it is not surprising that the timing of the main agricultural fire season peak and the more minor March–April peak we see both agree with the seasonality of cropland fires in Europe discussed by Korontzi et al. (2006).

The agricultural fire season in the Middle East (MIDE) lasts from May to November (Fig. 3). Yevich and Logan (2003) point out that farmers in Turkey, which has a high cropland fraction (Fig. 4), traditionally burn crop residues in the field. The winter wheat crop in the Middle East is planted from September to December and harvested from April to August (Leff et al., 2004), which is entirely consistent with our results. Namely, agricultural fires begin in June (after harvest) and end by November (before planting;

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fig. 3). Burning practices associated with pasture in other parts of MIDE (e.g., the Arabian Peninsula and the Mediterranean coast of Africa; Fig. 4) have not been well characterized in the literature.

The non-agricultural fire season in MIDE lasts from approximately March through October with a broad peak around May–June, corresponding to elevated levels of flammability and lightning frequency (Fig. 3).

4 Conclusions

Agricultural practices around the world have resulted in differences between the fire seasons associated with agriculture and those not associated with agriculture; we provide evidence for such differences in nearly every inhabited region on Earth (Fig. 3). We show that it is possible, using a concise algorithm, to decouple the agricultural and non-agricultural fire seasons by analyzing land-use data in conjunction with fire observations. Although we focused our discussion on burned area from GFED (Giglio et al., 2010), similar results for the fire count datasets from NASA MODIS (Justice et al., 2002; Giglio et al., 2006a) illustrate the flexibility and consistency of our method.

The method we used to separate the agricultural and non-agricultural fire seasons can be generalized to include additional land-use data. For example, deforestation is an important driver of fire in some regions (Morton et al., 2008); our method could be used to parameterize global fire models for this and significantly improve estimates of the timing and extent of fire there. Our method could also be used to distinguish between cropland versus pasture fire, providing an added dimension with respect to agricultural burning practices.

We perform our analysis in terms of broad study regions defined by past studies (e.g., Giglio et al., 2006b), but examination of the results suggests that the regions are often too large and simply defined. This leads to a mix of biomes and agricultural practices that increase the complexity of physical interpretation. The development of study regions for analyses of global fire patterns that are dependent on underlying

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



land cover and land use may be a simple but critical step in advancing the modeling of global fires beyond dependence of seasonality on climate conditions (e.g., Pechony and Shindell, 2009; Kloster et al., 2010; Thonicke et al., 2010) to include dependence on well-documented cultural burning practices (e.g., Yevich and Logan, 2003; Vigilante et al., 2004; Laris and Wardell, 2006; McCarty et al., 2009).

Our results here prove that fires used for management of agriculture and fires on non-agricultural land can be distinguished in observational time series of overall burning, especially if regional differences are considered to account for unique cultural practices. This suggests that seasonalities of different fire types in large-scale models can be parameterized by and/or tested against real patterns. Incorporating such anthropogenic influences into regional and global models of fire and vegetation will advance the scientific understanding of fire in the Earth system, both in the past and the future.

Supplementary material related to this article is available online at:

<http://www.biogeosciences-discuss.net/9/5551/2012/>

[bgd-9-5551-2012-supplement.pdf](#).

Acknowledgements. The authors were supported by the Carbon Mitigation Initiative and the Cooperative Institute for Climate Science at Princeton University.

References

- Aldersley, A., Murray, S. J., and Cornell, S. E.: Global and regional analysis of climate and human drivers of wildfire, *Sci. Total Environ.*, 409, 3472–3481, 2011.
- Arora, V. and Boer, G.: Fire as an interactive component of dynamic vegetation models, *J. Geophys. Res.*, 110, 1–20, 2005.
- Christian, H. J., Blakeslee, R. J., Boccippio, D. J., Boeck, W. L., Buechler, D. E., Driscoll, K. T., Goodman, S. J., Hall, J. M., Koshak, W. J., Mach, D. M., and Stewart, M. F.: Global fre-

BGD

9, 5551–5577, 2012

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



quency and distribution of lightning as observed from space by the optical transient detector, *J. Geophys. Res.*, 108, 4005, doi:10.1029/2002JD002347, 2003.

Giglio, L., Csiszar, I., and Justice, C. O.: Global distribution and seasonality of active fires as observed with the Terra and Aqua MODIS sensors, *J. Geophys. Res.*, 111, G02016, doi:10.1029/2005JG000142, 2006a.

Giglio, L., van der Werf, G. R., Randerson, J. T., Collatz, G. J., and Kasibhatla, P.: Global estimation of burned area using MODIS active fire observations, *Atmos. Chem. Phys.*, 6, 957–974, doi:10.5194/acp-6-957-2006, 2006b.

Giglio, L., Loboda, T., Roy, D. P., Quayle, B., and Justice, C. O.: An active-fire based burned area mapping algorithm for the MODIS sensor, *Remote Sens. Environ.*, 113, 408–420, doi:10.1016/j.rse.2008.10.006, 2009.

Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries, R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire products, *Biogeosciences*, 7, 1171–1186, doi:10.5194/bg-7-1171-2010, 2010.

Hansen, M., Defries, R. S., Townshend, J. R. G., and Sohlberg, R.: Global land cover classification at 1 km spatial resolution using a classification tree approach, *Int. J. Remote Sens.*, 21, 1331–1364, 2000.

Huffman, G. J., Adler, R. F., Bolvin, D. T., and Gu, G.: Improving the global precipitation record: GPCP Version 2.1, *Geophys. Res. Lett.*, 36, L17808, doi:10.1029/2009GL040000, 2009.

Justice, C. O., Giglio, L., Korontzi, S., Owens, J., Morrisette, J. T., Roy, D., Descloitres, J., Alleaume, S., Petitcolin, F., and Kaufman, Y.: The MODIS fire products, *Remote Sens. Environ.*, 83, 244–262, 2002.

Kalnay, E., Kanamitsu, M., Kitsler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–470, 1996.

Klein Goldewijk, K., Beusen, A., van Drecht, G., and de Vos, M.: The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years, *Global Ecol. Biogeogr.*, 20, 73–86, 2011.

Kloster, S., Mahowald, N. M., Randerson, J. T., Thornton, P. E., Hoffman, F. M., Levis, S., Lawrence, P. J., Feddes, J. J., Oleson, K. W., and Lawrence, D. M.: Fire dynamics during

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the 20th century simulated by the Community Land Model, *Biogeosciences*, 7, 1877–1902, doi:10.5194/bg-7-1877-2010, 2010.

Korontzi, S., McCarty, J., Loboda, T., Kumar, S., and Justice, C.: Global distribution of agricultural fire in croplands from 3 years of Moderate Resolution Imaging Spectroradiometer (MODIS) data, *Global Biogeochem. Cy.*, 20, GB2021, doi:10.1029/2005GB002529, 2006.

Laris, P. and Wardell, D. A.: Good, bad or “necessary evil?” Reinterpreting the colonial burning experiments in the savanna landscapes of West Africa, *Geogr. J.*, 172, 271–290, 2006.

Le Page, Y., Oom, D., Silva, J. M. N., Jönsson, P., and Pereira, J. M. C.: Seasonality of vegetation fires as modified by human action: observing the deviation from eco-climatic fire regimes, *Global Ecol. Biogeogr.*, 19, 575–588, 2010.

Leff, B., Ramankutty, N., and Foley, J. A.: Geographic distribution of major crops across the world, *Global Biogeochem. Cy.*, 18, GB1009, doi:10.1029/2003GB002108, 2004.

McCarty, J. L., Korontzi, S., Justice, C. O., and Loboda, T.: The spatial and temporal distribution of crop residue burning in the contiguous United States, *Sci. Total Environ.*, 407, 5701–5712, doi:10.1016/j.scitotenv.2009.07.009, 2009.

Morton, D. C., Defries, R. S., Randerson, J. T., Giglio, L., Schroeder, W., and van der Werf, G. R.: Agricultural intensification increases deforestation fire activity in Amazonia, *Glob. Change Biol.*, 14, 2262–2275, doi:10.1111/j.1365-2486.2008.01652.x, 2008.

Pechony, O. and Shindell, D. T.: Fire parameterization on a global scale, *J. Geophys. Res.*, 114, D16116, doi:10.1029/2009JD011927, 2009.

Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L., and Carmona-Moreno, C.: The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model, *Biogeosciences*, 7, 1991–2011, doi:10.5194/bg-7-1991-2010, 2010.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Vigilante, T., Bowman, D. M. J. S., Fisher, R., Russell-Smith, J., and Yates, C.: Contemporary landscape burning patterns in the far North Kimberly region of North-West Australia: human influences and environmental determinants, *J. Biogeogr.*, 31, 1317–1333, 2004.

Yevich, R. and Logan, J. A.: An assessment of biofuel use and burning of agricultural waste in the developing world, *Global Biogeochem. Cy.*, 17, 1095, doi:doi:10.1029/2002GB001952, 2003.

BGD

9, 5551–5577, 2012

**Separating
agricultural and
non-agricultural fire**

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Separating
agricultural and
non-agricultural fire**

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 1.** Descriptions of the regions.

Short name	Full name
BONA	Boreal North America
TENA	Temperate North America
CEAM	Central America and the Caribbean
NHSA	Northern Hemisphere South America
SHSA	Southern Hemisphere South America
EURO	Europe
MIDE	Middle East and Northern Africa
NHAF	Northern Hemisphere Africa
SHAF	Southern Hemisphere Africa
BOAS	Boreal Asia
CEAS	Central Asia
SEAS	Southeast Asia
EQAS	Equatorial Asia
AUST	Australia and New Zealand

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Fraction of the total number of grid cells in a region (N) that are predominantly non-agricultural vegetation (g_n) and predominantly agricultural vegetation (g_a).

Region	g_n	g_a	N
BONA	0.87	0.07	2651
TENA	0.24	0.30	2743
CEAM	0.17	0.14	928
NHSA	0.45	0.01	733
SHSA	0.35	0.10	4084
EURO	0.18	0.05	1898
MIDE	0.25	0.19	1313
NHAF	0.19	0.26	2630
SHAF	0.21	0.14	2858
BOAS	0.83	0.01	5673
CEAS	0.08	0.36	5172
SEAS	0.26	0.10	2050
EQAS	0.58	0.01	1032
AUST	0.05	0.05	2340

Separating agricultural and non-agricultural fire

B. I. Magi et al.




Fig. 1. Map of the regions in this study, after Giglio et al. (2006b). The short names are described in more detail in Table 1.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Separating agricultural and non-agricultural fire

B. I. Magi et al.




Fig. 2. Burned area from GFED (green) and fire counts from Terra MODIS (red) and Aqua MODIS (blue) for the regions in Fig. 1, normalized to the month of maximum burned area or fire counts. The x-axis shows months, while the y-axis shows the scaled burned area and fire counts. The numbers in parentheses are the percent contributions of the total mean annual fire counts or burned area in the region to the global total mean annual fire counts or burned area, following the same color scheme as the lines in the plot.

Separating agricultural and non-agricultural fire

B. I. Magi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 3. Agricultural (\bar{F}_a , green) and non-agricultural (\bar{F}_n , purple) fire seasonality derived from GFED for the regions in Fig. 1. The x-axis shows months and the y-axis shows \bar{F}_a and \bar{F}_n normalized to the maximum. The error bars indicate the interannual variability in \bar{F}_a and \bar{F}_n from 2000–2009. The dark gray line is the normalized flammability index, and the lighter gray line is the normalized lightning frequency.

Separating agricultural and non-agricultural fire

B. I. Magi et al.




Fig. 4. (a) Agricultural fraction of land area. (b) Composition of agricultural land, cropland vs. pasture. The figures show the average of 2000 and 2005 land-use distributions from the HYDE 3.1 data set. For both subfigures, gray denotes grid cells without agriculture.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

